

(19)

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 0 779 363 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

18.06.1997 Bulletin 1997/25

(51) Int Cl.⁶: C12N 15/82, C12N 15/54,
C12N 9/10, C08B 30/00,
C12N 5/10, A01H 5/00

(21) Application number: 96309004.8

(22) Date of filing: 11.12.1996

(84) Designated Contracting States:

AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE

- Marshall, Jacqueline
Warwick CV34 5SA (GB)
- Edwards, Elizabeth Ann
Norwich NR9 3DB (GB)
- Martin, Catherine Rosemary
Norwich NR15 1JW (GB)

(30) Priority: 12.12.1995 GB 9525353

(71) Applicant: National Starch and Chemical
Investment Holding Corporation
Wilmington, Delaware 19809 (US)

(74) Representative: Lipscombe, Martin John et al
Keith W Nash & Co,
Pearl Assurance House,
90-92 Regent Street
Cambridge CB2 1DP (GB)


(72) Inventors:

- Smith, Alison Mary
Norwich NR2 2BB (GB)


(54) Improvements in or relating to soluble starch synthase

(57) Disclosed is altered starch extracted from a transformed potato plant or the progeny thereof, which starch, as extracted, has a viscosity onset temperature as determined by differential scanning calorimetry which is reduced by at least 5°C compared to starch extracted from equivalent, non-transformed plants, together with a nucleic acid sequence comprising at least 200bp and exhibiting at least 80% sequence identity with the corresponding region of the DNA sequence shown in Figure 6, operably linked in the sense or anti-sense orientation to a promoter operable in a plant, and host cells and plants comprising the sequence.

A
pPRAT4



B
pPATRAT



2435 bp promoter
1.0 kb polyA
1.0 kb polyA
EcoRI
XbaI
2435 bp promoter
1.0 kb polyA
1.0 kb polyA
EcoRI
XbaI

EP 0 779 363 A2

Description

Field of the Invention

5 This invention relates, *inter alia*, to a soluble enzyme, obtainable from potato tubers, having starch synthase activity, to nucleic acid sequences encoding the same, to constructs and transgenic plants comprising the nucleic acid sequences, to a method of altering the starch composition of a plant, and to altered starch obtainable from a transgenic plant.

10 Background of the Invention

In the storage organs of most species of plants multiple forms of both granule-bound and soluble starch synthases have been found (for review, see Smith & Martin 1993, In: Biosynthesis and manipulation of plant products (D. Grierson, Ed.) Blackie Academic and Professional (Glasgow), pp1-54). In most cases it is not known whether these forms are
 15 distinct gene products and, for the most part, what their detailed functions are. The exception to this is in the case of a widely-distributed and highly conserved class of granule-bound starch synthases of approximately 60 kDa, which are collectively referred to as granule-bound starch synthase I (GBSS I; Martin & Smith, 1995 Plant Cell 7, 971-985). Experiments with the waxy and amf mutants of cereals and potatoes respectively (Macdonald & Preiss, 1985 Plant Physiol. 78, 849-852 1985; Hovenkamp-Hermelink *et al.*, 1987 Theor. Appl. Genet. 7, 217-221) and antisense potato
 20 plants (Visser *et al.*, 1991 Mol. Gen. Genet. 22, 289-296; Kuipers *et al.*, 1994 Plant Cell 6, 43-52) have shown that when the level of GBSS I protein is reduced, the ratio of amylose to amylopectin in the starch is also reduced. Where GBSS I is absent, the starch contains only amylopectin. This suggests that GBSS I is responsible for amylose synthesis.

However, the detailed functions of other isoforms of starch synthase are as yet unknown. In general, in conjunction with starch branching enzyme, they must be responsible for amylopectin synthesis but it is unknown whether different
 25 isoforms make different contributions to its structure. The first step in trying to understand the functions of these starch synthases is to characterise all of the isoforms in one organ. A few isoforms of starch synthase, other than GBSS I, have been identified at a detailed biochemical and molecular level in pea (Smith, 1990 Planta 182, 599-604; Denyer & Smith 1992 Planta 186, 609-617, Dry *et al.* 1992 Planta 186, 609-617) and rice (Baba *et al.*, 1993 Plant Physiol. 103, 565-573), and at a detailed biochemical level in maize (Mu *et al.*, 1994 Plant J. 6, 151-159) and wheat (Denyer
 30 *et al.*, 1995 Planta 196, 256-265). However only in the case of pea and maize has the quantitative importance of the isoforms been estimated. A complete picture of the role and importance of all the isoforms of starch synthase is not available for any other storage organ.

Carbohydrate metabolism and starch synthesis has been extensively studied in potato tuber (Hajirezaei *et al.*, 1993 Planta 192, 16-30; Geigenberger & Stitt 1993 Planta 189, 329-339; Geigenberger *et al.*, 1994 Planta 193, 486-493;
 35 Sonnewald *et al.*, 1994 Plant Cell Environ. 17, 649-658) and this organ has great potential as a source of commercially important starches created through genetic manipulation (Shewmaker & Stalker 1992 Plant Physiol. 100, 1083-1086; Visser & Jacobsen 1993 Trends Biotech. 11, 63-68; Muller-Rober & Koßmann, 1994 Plant Cell Environ. 17, 601-613). One of the major gaps in understanding starch synthesis in this organ and hence in the ability to manipulate its starch in useful ways, is the nature of its starch synthases.

In potato, until recently, only two starch synthases have been characterised in any great detail: GBSS I and GBSS II. GBSS I is exclusively granule-bound, it has a molecular weight of 59 kDa. The gene has been cloned and its predicted amino acid sequence is very similar to that of the waxy gene product in cereals (Vos-Scheperkeuter *et al.*, 1986 Plant
 40 Physiol. 82, 411-416; van der Leij *et al.*, 1991 Mol. Gen. Genet. 228, 240-248). GBSS II has an apparent molecular weight, as judged by SDS-PAGE, of 92 kDa and it is both bound into the starch granule and present as a soluble form. Its predicted amino acid sequence (having an expected molecular weight of 80 kDa) is similar to GBSS II in pea embryos, an isoform which accounts for 60-70 % of the soluble starch synthase activity of the pea embryo (Denyer & Smith 1992
 45 cited above). However, GBSS II accounts for only approximately 10-15% of the total soluble starch synthase activity in potato tubers (Edwards *et al.*, 1995 Plant J. 8, 283-294).

There have been several reported characterisations of the starch synthases found in the soluble fraction of potato
 50 tubers (Frydman & Cardini 1966 Arch. Biochem. Biophys. 116, 9-18; Catz *et al.*, 1989 An. Asoc. Quim. Argent. 77, 47-51) and a few attempts have been made to purify the major soluble starch synthases (Hawker *et al.*, 1972. Phytochem. 11, 1278-1293; Baba *et al.*, 1990 Phytochem. 29, 719-723; Ponstein 1990. Starch synthesis in potato tubers. Ph. D. Thesis, State University Groningen, The Netherlands). These reports disagree on both the number of soluble starch synthases and their molecular weights. The quantitative contribution of the putative forms is not known, and
 55 where multiple forms are postulated, it is not known whether they are independent gene products.

After the priority date of the present application, two publications have been made which provide information about a further starch synthase found in potato. One of these publications is by the present inventors (Marshall *et al.*, 1996 The Plant Cell 8, 1121-1135). The other publication is PCT patent application WO 96/15248 (published 23rd May 1996),

in the name of Institut Für Genbiologische Forschung Berlin GMBH. The PCT application includes the European Patent Office in the list of designated territories.

WO 96/15248 discloses the nucleotide sequence of a full length cDNA clone ("SSSA") said to encode an isoform of soluble starch synthase enzyme from potato, together with the predicted amino acid sequence of the enzyme. The application further discloses the use of a 1.2kb portion of the cDNA clone, operably linked in the antisense orientation to the CaMV 35S promoter, to transform potato plants. In addition WO 96/15248 discloses the sequence of a cDNA clone ("SSSB") said to encode a second isoform of the potato soluble starch synthase. Similarly, a portion (1.8kb) of this sequence was introduced into potato plants in the antisense orientation.

In fact, the present inventors have found that the nucleotide sequence disclosed in WO 96/15248 contains an error, causing a frame shift, such that most of the predicted amino acid sequence is incorrect.

It was found that the transformed plants disclosed in WO 96/15248 had reduced enzyme activity. Starch obtained from the tubers of the transformed plants was found to have altered properties compared to starch obtained from control wild type plants. It was stated that the starch from the transformed plants exhibited a lower viscosity onset temperature than starch from control plants. (By way of explanation, when aqueous suspensions of starch granules are heated, the granules swell and absorb water, in a process known as gelatinisation. A number of techniques are available for the analysis of gelatinisation, a particularly convenient method being differential scanning calorimetry or the viscoamylograph, in which the viscosity of a stirred starch suspension is monitored under a defined temperature/time regime. Such analysis typically shows a particular temperature, the "viscosity onset temperature", at which the process of gelatinisation begins and which causes a marked increase in viscosity of the starch suspension).

In a few instances, the transformed plants disclosed in WO 96/15248 gave rise to starch in which the "Verkleisterungstemperatur" (equivalent to the viscosity onset temperature, V) was 2 to 3°C lower compared to starch from equivalent, but untransformed, plants. However, it is apparent that the results of subsequent experiments (described in example 13 in the document) gave a value of V for starch from control plants which was lower than that found for starch from transformed plants in previous experiments. Accordingly, the person skilled in the art is not able to deduce that starch from the transformed plants described in WO 96/15248 displayed a viscosity onset temperature which was consistently significantly lower than that of control plants.

Summary of the Invention

In a first aspect the invention provides a polypeptide obtainable from a soluble extract of potato tubers and having starch synthase activity, in substantially isolated form.

Typically the polypeptide will have an apparent molecular weight, as judged by SDS-PAGE, in the range 100-140 kDa, or will be a functional equivalent of such a polypeptide. More particularly, the polypeptide may have an apparent molecular weight of 140, 120 or 110 kDa. Particular functional equivalents envisaged are breakdown products of the polypeptide, which seem to occur naturally. Another particular functional equivalent is the polypeptide obtainable from developing tubers of the Desiree cultivar, which polypeptide has an apparent molecular weight, as judged by SDS-PAGE, of 100 kDa. Typically the polypeptide will comprise the amino acid sequence shown in Figure 6.

In another aspect the invention provides a nucleic acid sequence directing the expression of at least a portion of one of the polypeptides defined above. Preferably the sequence comprises at least 200-300bp, more preferably at least 300-600bp, and most preferably in excess of 600bp. Typically the nucleic acid sequence will comprise the nucleotide sequence shown in Figure 6, although those skilled in the art will appreciate that, due to the degeneracy of the genetic code, a nucleotide sequence substantially different to that shown in Figure 6 may encode a polypeptide having substantially the same amino acid sequence as that shown in Figure 6. Such nucleic acid sequences are to be considered as functional equivalents and thus fall within the scope of the present invention. Other functional equivalents are those nucleic acid sequences which are not substantially different and which may hybridise, under standard laboratory hybridisation conditions, to either strand of the nucleotide sequence shown in Figure 6.

Comparison with known starch synthase sequences, with the benefit of the disclosure herein, will enable those skilled in the art to identify regions of the sequence shown in Figure 6 which are not evolutionarily conserved, and so more amenable to alteration (e.g. addition, deletion or substitutions), whilst retaining functional equivalence.

Desirably such functional equivalents will possess at least 80% sequence identity, preferably at least 85% sequence identity, and more preferably at least 90% sequence identity with the nucleotide sequence shown in Figure 6. Desirably the nucleotide sequence of the invention, or a functional equivalent sequence will, when introduced into a suitable plant in a suitable manner (known to those skilled in the art), alter the synthesis of starch in the plant.

For the purposes of the present specification, the sequences encoding polypeptides with starch synthase activity, or portions of such sequences, disclosed in WO 96/15248 are not considered as functional equivalents of the sequence shown in Figure 6.

In a particular embodiment, the invention provides a nucleic acid sequence comprising at least 200 bp and exhibiting at least 80% sequence identity with the corresponding region of the DNA sequence shown in Figure 6, operably

linked in the sense sense orientation to a promoter operable in a plant.

Those skilled in the art will readily be able to conduct a sequence alignment between the other sequence and that detailed in Figure 6. The % identity of the two sequences is to be compared in those regions which are aligned by readily available computer programs (e.g. MegAlign), which align corresponding regions of sequences. Advantageously the % identity between the two sequences will be at least 85%, preferably at least 90%, and the corresponding region of the sequence shown in Figure 6 may comprise a 5' and/or a 3' untranslated region ("UTR") and/or a translated region.

Thus, in another aspect the invention provides a nucleic acid construct (typically DNA) comprising the nucleic acid sequence of the invention in operable linkage to a promoter active in a plant. The nucleic acid sequence may be operably linked to the promoter in either the sense or the anti-sense orientation. Anti-sense methods are well known in altering one or more characteristics of a plant into which the anti-sense sequence is inserted (see for example EP-A-0 458 367, EP-B-0 240 208 and US 5, 107, 065). Similarly, "sense suppression" is a method which is becoming increasingly widely adopted and documented (for a review, see Matzke & Matzke 1995 Plant Physiology 107, 679-685). Either approach could be used with the nucleic acid sequence of the invention, so as to alter one or more characteristics of a plant into which the sequence was introduced. Those skilled in the art will be aware that anti-sense inhibition or sense suppression may be achieved by the use of 5' or 3' non-translated portions of the relevant gene, or use of coding portions of the gene, or any combination thereof.

It is believed that antisense methods are mainly operable by the production of antisense mRNA which hybridises to the sense mRNA, preventing its translation into functional polypeptide, possibly by causing the hybrid RNA to be degraded (e.g. Sheehy *et al.*, 1988 PNAS 85, 8805-8809; Van der Krol *et al.*, Mol. Gen. Genet. 220, 204-212). Sense suppression also requires homology between the introduced sequence and the target gene, but the exact mechanism is unclear. It is apparent however that, in relation to both antisense and sense suppression, neither a full length nucleotide sequence, nor a "native" sequence is essential. Preferably the nucleic acid sequence used in the method will comprise at least 200-300bp, more preferably at least 300-600bp, of the full length sequence, but by simple trial and error other fragments (smaller or larger) may be found which are functional in altering the characteristics of the plant. It is also known that untranslated portions of sequence can suffice to inhibit expression of the homologous gene - coding portions may be present within the introduced sequence, but they do not appear to be essential under all circumstances.

The invention further provides a host cell into which has been introduced a nucleic acid sequence in accordance with the invention defined above. Typically the host cell will be a plant cell, and conveniently the sequence is introduced in a nucleic acid construct and subsequently integrated into the host cell genome.

In a further aspect the invention provides a plant or part thereof (e.g. plant cell), into which has been introduced a sequence in accordance with the invention, or the progeny of such a plant or part thereof. Desirably the plant or part thereof into which the sequence is introduced, will comprise a natural gene which shares sequence homology with the introduced sequence. In preferred embodiments the introduced sequence will exhibit at least 70% homology with a starch synthase gene naturally present in the plant or part thereof, although the level of homology may be increased with advantage, such that the expression of the gene product of the naturally present gene in the plant is substantially inhibited. Conveniently the sequence of the invention will be introduced as part of a nucleic acid construct, as described above. Typically the plant will be one of commercial significance, such as one of the following: potato, tomato, rice, wheat, pea, cassava, sweet potato, barley, oat and maize.

Those skilled in the art will appreciate that introduction of the nucleic acid sequence of the invention into a plant might alter the starch composition thereof. In another aspect therefore the invention provides altered starch extracted from a plant into which has been introduced the nucleic acid sequence of the invention, or altered starch extracted from the progeny of such a plant. The invention also provides a method of altering one or more characteristics of a plant, comprising introducing into the plant a nucleic acid sequence in accordance with the invention.

In particular the invention provides altered starch extracted from a transformed potato plant or the progeny thereof, which starch, as extracted, has a viscosity onset temperature as determined by differential scanning calorimetry which is reduced by at least 5°C compared to starch extracted from equivalent, non-transformed plants. Preferably the viscosity onset temperature is reduced by at least 7°C. In another embodiment the invention provides altered starch extracted from a transformed potato plant or the progeny thereof, which starch, as extracted, has a viscosity onset temperature as determined by differential scanning calorimetry, of less than 60°C, preferably less than 55°C.

The starch defined above will typically also exhibit (as extracted) a reduced endotherm peak temperature, as determined by differential scanning calorimetry, compared to starch extracted from equivalent, non-transformed plants. Desirably the endotherm peak temperature will be reduced by at least 5°C and/or will be less than 59°C. The inventors have found that such properties as those defined above may be embodied in potato starch having a substantially normal amylose content (i.e. around 25 - 30% amylose).

Starch can be modified in various ways (e.g. chemical cross-linking, derivatisation, partial hydrolysis) after it has been extracted from a plant source, which modifications can affect the physical properties, especially the pasting properties, of the starch. Hence, use of the term "as extracted" is intended to signify that the starch is analysed without

undergoing such modifications as can alter the pasting properties thereof.

"Equivalent, non-transformed" plants are those plants which have substantially identical genotypes to the plants of the invention, with the exception of the introduced nucleic acid sequence present in the transformed plants of the invention.

5 The invention will now be further described by way of illustrative example and with reference to the accompanying drawings, in which:

10 Figure 1 shows the elution profile of starch synthase from developing Desiree potato tubers on a first Mono Q™ anion-exchange column. Partially purified starch synthase, after DEAE-Sepharose and Blue Sepharose chromatography, was applied to a 1ml Mono Q™ column at pH 7.5. The enzyme was eluted with a 25ml gradient of 0-450 mM KCl at 0.5 ml.min⁻¹. Samples (20 µl) of each 1 ml fraction were assayed for starch synthase activity (●), and absorbance at 280 nm (-):

15 Figure 2 shows the activity and protein in fractions of purified starch synthase from a second Mono Q™ column of peak I and peak II. Top panels show SDS-PAGE of fractions containing starch synthase activity. Each track contains 10 µl of fraction. Bottom panels show starch synthase activity in 20 µl samples from each 0.5 ml fraction:

20 Figure 3 shows the cross-reaction of antiserum to SSS to the purified starch synthases from mature Estima tubers and to extracts, soluble and granule-bound, from mature Estima and developing Desiree tubers. Samples (10 µl of purified soluble starch synthase, 20 µl of partially purified soluble starch synthase, 20 µl soluble extract and 20 µl of supernatant from granule-bound proteins) were subjected to SDS-PAGE and blotted, and then the blots were probed with antiserum to SSS, 1/2500 dilution. (1) purified preparation of starch synthase proteins from mature Estima. (2) Partially purified soluble starch synthase from mature Estima tubers. (3) Starch-granule-bound proteins from mature Estima tubers. (4) Soluble extract from developing Desiree tubers. (5) Starch-granule-bound proteins from developing Desiree tubers. Sizes of proteins were estimated from molecular weight standards on the same gels, and are indicated in kDa:

25 Figure 4 shows the immunoprecipitation of starch synthase activity in soluble extract from developing Desiree tubers with antiserum to SSS. Soluble extract was incubated with increasing volumes of pre-immune serum (○) and antiserum (●), as described in Materials and Methods (below). After centrifugation the supernatant was assayed for starch synthase activity. Starch synthase activity is expressed as a percentage of activity of incubations containing 20 g.L⁻¹ BSA in PBS. Values are from two separate experiments with the line joining the means:

30 Figure 5 shows native polyacrylamide gel electrophoresis of soluble extract from developing Desiree tubers stained for starch synthase activity. Soluble extract was incubated (as described in Materials and Methods) in the presence of (1) 20 g.L⁻¹ BSA in PBS; (2) pre-immune serum, 1/1000 dilution; (3) antiserum to SSS, 1/1000 dilution; and (4) antiserum to the GBSS II from pea embryo. After centrifugation, the supernatant was mixed 5:1 with 2 g.L⁻¹ bromophenol blue in 500 ml.L⁻¹ glycerol and 40 µl was loaded onto the gel. The bands of starch synthase activity are indicated by arrows:

35 Figure 6 shows the DNA sequence of a cDNA clone for potato soluble starch synthase. The amino acid sequence of the encoded polypeptide is shown below in the single letter code. The ADP-glucose binding domain is boxed and the sequences identified by protein sequencing are underlined: and

40 Figure 7 shows a schematic representation of A) plasmid pRAT4 and B) plasmid pPATRAT.

45 **EXAMPLES**

50 **Example 1**

In this example are presented data on the identification and purification to homogeneity of the major isoform of soluble starch synthase from potato tuber.

55 **MATERIALS AND METHODS**

Plant material.

Potato tubers (*Solanum tuberosum* L.) of cultivars Desiree (developing) or Estima (mature) were used. Desiree

tubers were grown in soil based compost (25 cm diameter) in a greenhouse with minimum temperature of 12°C and supplementary lighting in winter, and were freshly harvested prior to experiments from actively-growing plants. Estima tubers were bought locally.

5 Purification of soluble starch synthase.

(A) Small scale. The extraction and subsequent purification were carried out at 4°C. Approximately 500g of Desiree potato tubers were chopped into small pieces and homogenised in an electric blender with 25g polyvinylpyrrolidone (PVPP) and 500 ml of ice-cold medium A containing 100 mM Tris-HCl (pH 7.5), 10 mM ethylenediamine-tetraacetic acid (EDTA), 5 mM dithiothreitol (DTT), 1 g.L⁻¹ sodium metabisulphite, 0.5 mg.L⁻¹ leupeptin, 0.7 mg.L⁻¹ pepstatin A and 50 ml.L⁻¹ glycerol. The homogenate was passed through two layers of muslin and the filtrate was centrifuged at 10,000g for 10 min. The supernatant was brought to 40% saturation with powdered (NH₄)₂SO₄. The precipitate was collected by centrifugation (15,000g for 15 min), re-dissolved in a minimal volume of medium A and dialysed twice, each time against 1 L of medium A for 1 h.

The dialysed extract was applied, at a flow rate of 4 ml.min⁻¹, to a column (5 cm internal diameter "i.d.", 10 cm long) of diethylaminoethyl (DEAE)-Sephacel™ (Pharmacia, Uppsala, Sweden), equilibrated with medium A. The column was washed with 500 ml of medium A followed by a 250-ml gradient of 0-1 M KCl in the same medium. Fractions of 10 ml were collected and assayed for starch synthase activity. The eight to ten fractions containing the highest activity were pooled and dialysed twice, each time against 1 L of medium B containing 50 mM Tris-HCl (pH 7.5), 1 mM EDTA, 1 mM DTT, 0.5 mg.L⁻¹ leupeptin, 0.7 mg.L⁻¹ pepstatin A and 50 ml.L⁻¹ glycerol for 1 h.

The dialysed extract was applied, at a flow rate of 1 ml.min⁻¹, to a column (1.6 cm i.d., 16 cm long), of Blue Sepharose, equilibrated with medium B. The column was washed with 100 ml medium B followed by a 100ml gradient of 0-1 M KCl in the same medium. Fractions of 0.5 ml were collected and assayed for starch synthase activity. The ten fractions with the highest activity were pooled and dialysed twice, each time against 1 L of medium B for 1 h.

The dialysed extract was applied, at a flow rate of 0.5 ml.min⁻¹, to a first 1-ml Mono Q™ column (Pharmacia), equilibrated with medium B. The column was washed with 25 ml of medium B, followed by a 25-ml gradient of 0-450 mM KCl in the same medium. Fractions of 0.5 ml were collected and assayed for starch synthase activity. The fractions from each of two peaks of starch synthase activity were pooled and purified separately as follows:

To the eluate of the first Mono Q™ column, an equal volume of 1 M sodium citrate in medium B was added. This was then applied, at a flow rate of 0.5 ml.min⁻¹, to a column (1.0 cm i.d., 4 cm long) of cyclohexa-amylose (CHA)-Sephacel (prepared according to Vretblad, 1974 FEBS Lett. 47, 86-89), equilibrated with 0.5 M sodium citrate in medium B. The column was washed with 20 ml medium B containing 0.5 M sodium citrate and the protein was eluted from the column with 30 ml of medium B containing no citrate. Fractions of 1 ml were collected and assayed for starch synthase activity. The fractions with starch synthase activity were pooled and dialysed overnight against 5 L of medium C containing 50 mM Tris-HCl (pH 8.0), 1 mM EDTA, 1 mM DTT, 0.5 mg.L⁻¹ Leupeptin, 0.7 mg.L⁻¹ pepstatin A and 50 ml.L⁻¹ glycerol.

The dialysed extract was applied, at a flow rate of 0.5 ml.min⁻¹, to a second 1ml Mono Q™ column equilibrated with medium C. The column was washed with 25 ml of medium C followed by a 25ml gradient of 0-450 mM KCl in the same medium. Fractions of 0.5 ml were collected and assayed for starch synthase activity.

(B) Large scale.

The procedures were as described above, with the following modifications. Five kg of Estima potato tubers were homogenised in 5 L medium A containing 250g PVPP, filtered through two layers of muslin and centrifuged at 10,000g for 10 min. Polyethylene glycol (PEG) 6000 at a concentration of 500 g.L⁻¹ in medium A was slowly added to the supernatant until the concentration of PEG was 100 g.L⁻¹. The precipitate was collected by centrifugation (15,000g for 20 min) and re-dissolved in a minimal volume of medium A.

The extract was mixed gently for 1 h with 900ml slurry of DEAE-Sephacel which had been equilibrated with medium A, then filtered and the filtrate discarded. The DEAE-Sephacel was washed with 2 L medium A then incubated for 1 hr in 500 ml medium A containing 400 mM KCl, filtered and washed with a further 500 ml medium A containing 400 mM KCl. The filtrates were combined and brought to 50% saturation with powdered (NH₄)₂SO₄. The precipitated proteins were collected by centrifugation (15,000g for 15 min), re-dissolved in a minimal volume of medium B and dialysed overnight against 5 L of medium B.

The dialysed sample was applied, at a flow rate of 2 ml.min⁻¹, to a Blue Sepharose column (5 cm i.d., 15 cm long) which had been equilibrated with medium B. The column was washed with 300 ml medium B followed by a 600ml gradient of 0-1 M KCl in the same medium, at a flow rate of 5 ml.min⁻¹. Fractions of 15 ml were collected and assayed for starch synthase activity. The ten fractions with the highest starch synthase activity were pooled and dialysed overnight against 5 L medium B.

The dialysed eluate was applied to a first 1ml Mono Q™ column, equilibrated with medium B, as described above, except that all the fractions containing starch synthase activity were pooled together.

The Mono Q™ eluate was applied to a CHA-Sepharose column as described above, except that the column was 1.0 cm i.d., 20 cm long. The column was washed with 50 ml medium B containing 0.5 M sodium citrate and eluted with 80 ml medium B without citrate. The fractions with starch synthase activity were pooled and dialysed overnight against 5 L of medium C.

The dialysed extract was applied to a second 1ml Mono Q™ column equilibrated with medium C, as described above. Fractions containing starch synthase activity were stored at -20°C.

10 Preparation of antibody.

The fractions containing starch synthase activity from 5 large-scale purifications were run on preparative sodium dodecyl sulphate (SDS)-polyacrylamide gels (as described below). The gel slices containing starch synthase proteins were electroeluted and the proteins dialysed against water, then freeze-dried. Protein (50 µg) was re-dissolved in 250 µl of phosphate-buffered saline (PBS), mixed with 250 µl Freund's complete adjuvant, and injected intramuscularly into a rat. Subsequent injections were of 75 µg protein dissolved in 250 µl PBS mixed with 250 µl Freund's incomplete adjuvant and were repeated at 14-day intervals. Serum was collected from 14 days after the third injection.

20 Assay of soluble starch synthase activity.

Soluble starch synthase activity was measured using the resin method as described in Jenner *et al.* (1994).

Preparation of crude soluble potato tuber extract.

25 Samples (0.5-2.0 g fresh weight) from either developing Desiree or mature Estima potato tuber were homogenised in 4 volumes of 50 mM Tris-HCl (pH 7.5), 1 mM EDTA, 1 mM DTT, 1 g.L⁻¹ sodium metabisulphite, 0.5 mg.L⁻¹ leupeptin, 0.7 mg.L⁻¹ pepstatin A and 50 ml.L⁻¹ glycerol, then centrifuged at 10,000g for 10 min. The supernatant is referred to as "soluble extract".

30 Partial purification of soluble starch synthase activity.

Crude soluble potato extract from mature Estima tubers (5-10 g fresh weight) was dialysed twice, each time against 1 L of buffer B for 1 hr. The dialysed extract was applied to a 1ml Mono Q™ column, equilibrated with medium B, as described above and the peak fraction of starch synthase activity (referred to as "partially purified soluble starch synthase") was stored at -20°C.

SDS-polyacrylamide gel electrophoresis (PAGE) and immunoblotting.

40 Protein samples were dialysed against distilled water then mixed 1:1 with double-strength sample buffer (Laemmli, U.K. (1970). Nature 227, 680-685) and boiled for 2 min immediately prior to application to gels. For granule-bound proteins, starch granules were washed twice in 20 g.L⁻¹ SDS at room temperature, boiled for 3 min at 100 mg.ml⁻¹ in sample buffer (Laemmli, U.K. (1970). Nature 227, 680-685) and then centrifuged at 10,000g for 10 min. The supernatant was applied to the gel.

45 Gels (10.2 cm long, 7.3 cm wide, 0.75 mm thick) were 75 g.L⁻¹ acrylamide (37.5:1 w/w acrylamide:bis-acrylamide) and 1 g.L⁻¹ SDS and were run according to Laemmli (1970). Immunoblots were prepared and developed according to Bhattacharyya *et al.*, (1990) Cell 60, 115-122. The nitrocellulose filters were either incubated with crude rat serum followed by alkaline phosphatase-conjugated goat anti-rat antiserum (Sigma, Poole, Dorset, UK) or the immunoglobulin fraction of rabbit serum to GBSS II from pea embryo (Smith (1990) Planta 182, 599-604), followed by alkaline phosphatase-conjugated goat anti-rabbit antiserum (Sigma).

Native polyacrylamide gel electrophoresis.

55 Gels (dimension as above, except 1mm thick) of 90 g.L⁻¹ acrylamide (37.5:1 w/w acrylamide:bis-acrylamide) were cast in 400 mM Tris-HCl (pH 8.6), 100 ml.L⁻¹ glycerol, 8 g.L⁻¹ glycogen and polymerised with 0.4 g.L⁻¹ ammonium persulphate and 0.2 ml.L⁻¹ N,N,N',N'-tetramethylethylenediamine (TEMED) and were overlaid with a stacking gel of 53 g.L⁻¹ acrylamide (37.5:1 w/w acrylamide:bis-acrylamide) cast in 155 mM Tris-HCl (pH 6.8), 98 ml.L⁻¹ glycerol, polymerised with 0.5 g.L⁻¹ ammonium persulphate and 0.2 ml.L⁻¹ TEMED. Soluble extracts were mixed 5:1 with 2 g.L⁻¹ bromophenol blue in 500 ml.L⁻¹ glycerol immediately prior to loading. Gels were run at 4°C, at 175 mV in 190 mM

glycine, 25 mM Tris

The gel was assayed for starch synthase activity as follows. The gel was washed twice, each time for 10 min in 20 ml 100 mM Bicine, 0.5 M sodium citrate (pH 8.5), 0.5 M EDTA and 100 ml.L⁻¹ glycerol at 4°C. The gel was incubated at room temperature for 20 hrs by gently shaking in wash medium containing 12 mM ADPG and 2 mM DTT. The buffer was removed and 1 ml of Lugol's iodine solution (3.3 g.L⁻¹ I₂ and 6.7 g.L⁻¹ KI, acidified with a few drops of 2M HCl) was added. After colour development, the gel was washed and stored in 70 ml.L⁻¹ acetic acid.

Immunoprecipitation.

Soluble extracts (100µl) were incubated with 0-20µl rat serum or 20µl of the immunoglobulin fraction of rabbit serum to GBSS II from pea embryo (Smith (1990) *Planta* 182, 599-604) for 1.5 h at room temperature on a rotating table. To the extract containing rat serum, 20µl polyclonal antiserum to rat IgG at 2.5 g.L⁻¹ specific antibody (Sigma) was added and incubated for a further 0.5 h. To both extracts, 50 µl Protein A-Sepharose at 60 g.L⁻¹ in 50 mM Tris-HCl (pH 7.5) was added and then incubated for 0.5 h, followed by centrifugation at 10,000g for 10 min. The supernatants were assayed for starch synthase activity. Controls contained bovine serum albumin at 20 g.L⁻¹ in PBS in place of serum.

Isolation of starch granules.

Purified starch was prepared from potato tubers as described by Edwards *et al.* (1995).

Measurement of protein.

Protein was assayed using the BioRad Protein Assay Dye Reagent (BioRad Munchen, Germany) with a standard curve of bovine serum albumin.

RESULTS

Purification of soluble starch synthases.

The soluble starch synthase activity from developing tubers of Desirée and mature tubers of Estima eluted from both DEAE-Sepharose and Blue Sepharose columns as a single peak of activity. However, subsequent chromatography on a Mono Q™ column at pH 7.5 separated two major peaks of starch synthase activity, designated peak I and peak II according to their elution order from the column (Figure 1). These two peaks of starch synthase activity were then purified separately by cyclohexa-amylose and Mono Q™ chromatography. A typical purification from developing Desirée tubers is shown in Table 1. The specific activity of peak I was 5.1 µmol.(mg protein)⁻¹.min⁻¹, a purification of 400-fold relative to the initial supernatant. The specific activity of peak II was 8.8 µmol.(mg protein)⁻¹.min⁻¹, a purification of 700-fold relative to the initial supernatant (Table 1).

Table 1 shows the purification of soluble starch synthase from developing potato tubers of Desirée. Fractions were prepared as described above. The values shown in the table are from a typical purification.

TABLE 1

FRACTION	TOTAL ACTIVITY (µmol glucose incorporated min ⁻¹)	ACTIVITY RECOVERED (%)	TOTAL PROTEIN (mg)	SPECIFIC ACTIVITY (µmol glucose incorporated min ⁻¹ . mg protein ⁻¹)
Initial Supernatant	28.9	100	2210.6	0.013
0 to 40% (NH ₄) ₂ SO ₄	17.1	61.0	1018.6	0.017
DEAE-Sepharose	7.51	26.8	45.1	0.166
Blue-Sepharose	4.59	16.4	10.4	0.441
Peak I				
Mono Q (pH 7.5)	0.95	3.4	1.70	0.56
Cyclohexa-amylose	0.53	1.9	0.20	2.65

TABLE 1 (continued)

FRACTION	TOTAL ACTIVITY (μmol glucose incorporated min^{-1})	ACTIVITY RECOVERED (%)	TOTAL PROTEIN (mg)	SPECIFIC ACTIVITY (μmol glucose incorporated min^{-1} mg protein $^{-1}$)
Peak I				
Mono Q (pH 8.0)	0.15	0.5	0.03	5.13
Peak II				
Mono Q (pH 7.5)	2.45	8.8	2.90	0.85
Cyclohexa-amylose	2.27	8.1	0.40	5.67
Mono Q (pH 8.0)	0.26	0.9	0.03	0.84

TABLE 2

INCUBATION	INHIBITION OF STARCH SYNTHASE ACTIVITY (%)
Pre-immune serum	0.3 ± 0.9
Antiserum to potato SSS	74 ± 4
Antiserum to pea GBSS II	9 ± 4
Antiserum to potato SSS + pea GBSS II	80 ± 8

Table 2 shows the immunoprecipitation of starch synthase activity in soluble extract from developing Desiree tubers. Soluble extract was incubated in the presence of antiserum (1/10 dilution of rat antiserum; or 1/5 dilution of the immunoglobulin fraction of rabbit serum to GBSS II from pea embryo; or 1/10 dilution of rat antiserum plus 1/5 dilution of the immunoglobulin fraction of rabbit serum to GBSS II), as described in Material and Methods. After centrifugation the supernatant was assayed for starch synthase activity. Values are percentage inhibition relative to controls in which BSA at 20 g.l^{-1} in PBS was substituted for serum. The values are the mean of four experiments \pm standard error.

Sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE) of the fractions from the second Mono QTM column for peak I showed that the distribution of a protein of 120 kDa matched the distribution of the starch synthase activity (Figure 2). Further chromatography on Mono QTM did not eliminate contaminating proteins. SDS-PAGE of the fractions from the final Mono QTM column for peak II showed that the distribution of the major protein of 110 kDa matched the starch synthase activity (Figure 2).

Antibodies raised to the 59 kDa starch-granule-bound protein (the GBSS I isoform) from pea embryo did not recognise any proteins from either peak I or peak II. Antibodies raised to the 77 kD GBSS II from pea embryo very weakly recognised the 120 and 110 kDa proteins from peak I and II respectively (data not shown).

Preparation of antibody.

In order to obtain sufficient protein for preparation of an antibody, peaks I and II from mature Estima tubers were combined and purified together in large-scale preparations (referred to as "soluble starch synthase", SSS). Both the 120- and 110 kDa proteins were excised and eluted from gels of the purified preparations and were injected into the same rat.

The antiserum to the SSS was used to probe blots of extracts from mature Estima and developing Desiree tubers (Figure 3). On all of the immunoblots, the pre-immune serum did not cross-react with any of the proteins. On immunoblots of the gels of the purified preparation of soluble starch synthase from mature Estima tubers, the antiserum recognised strongly the two proteins to which it was raised. The antiserum also recognised a minor protein of 140 kDa. On immunoblots of gels of partially purified soluble starch synthase from mature Estima tubers, the antiserum recognised proteins of 140 kDa and 120 kDa. On immunoblots of gels of starch-granule-bound proteins from mature Estima tubers, the antiserum recognised a protein of 140 kDa. There were some faint indications that a 120 kDa protein on the starch was also recognised. A protein of 140 kDa was recognised by the antiserum both in the soluble extracts and on starch granules of developing tubers of Desiree. The 120 kDa protein was very weakly detectable in soluble extracts from these tubers, which also contained a lower molecular weight protein (approximately 100 kDa) recognised

by the antiserum. This protein is not GBSS II since the rat antiserum did not recognise GBSS II on starch granules (data not shown).

Immunoprecipitation of starch synthase activity.

To discover whether the proteins recognised by the antiserum to SSS represent the major soluble starch synthases, the antiserum was used in immunoprecipitation experiments with soluble extracts from developing Desiree tubers.

Incubation of soluble extract with pre-immune serum from the rat did not affect soluble starch synthase activity, but the antiserum to SSS precipitated starch synthase activity (Figure 4). The maximum inhibition of starch synthase activity was approximately 75 % which was achieved by incubating with volumes greater than 2 μ l of antiserum. A small proportion of the remaining starch synthase activity can be accounted for by GBSS II (Table 2). When soluble extract is incubated with antiserum raised to GBSS II from pea embryo (which recognises GBSS II in potato, Edwards *et al.*, 1995), approximately 9% of the starch synthase activity is inhibited. When the potato extract is mixed with both antibodies, the starch synthase activity is reduced by approximately 80%.

Native polyacrylamide gel electrophoresis of soluble extracts of developing Desiree tubers revealed two major groups of bands which had starch synthase activity (Figure 5). We have previously shown through antisense and immunoprecipitation experiments that the lower group of bands is attributable to GBSS II. Tubers in which GBSS II protein has been severely reduced by antisense transformation lack the lower group of bands. When the soluble extract was immunoprecipitated with antiserum to GBSS II from pea embryo and the supernatant subjected to native PAGE, the lower bands were missing (Edwards *et al.*, 1995 cited previously). Immunoprecipitation of soluble extract from developing Desiree tubers with rat antiserum to SSS shows that the upper group of bands is attributable to these starch synthases. When the supernatant from the immunoprecipitation experiment was subjected to native PAGE, the upper group of bands was missing but the lower group was unaffected. The pre-immune serum from rat had no effect on the bands of starch synthase activity.

DISCUSSION

The inventors have purified two proteins with starch synthase activity from the soluble fraction from mature Estima tubers, with molecular weights of 110 and 120 kDa respectively. Immunoblots show that the antiserum raised to these purified proteins (soluble starch synthases, SSS) recognises the proteins to which it was raised, and that it also recognises a higher molecular weight protein of 140 kDa in the purified preparation (Figure 3). The 140 kDa protein is in soluble extracts and on starch granules of both mature Estima and developing Desiree tubers, whereas the 120 kDa protein is either barely or not detectable in tubers, and the 110 kDa protein is not detectable at all. This strongly suggests that the two starch synthase proteins to which antibodies have been raised may both be active breakdown products of the larger 140 kDa protein, although the 140 kDa polypeptide might simply be an immunologically cross-reactive entity. Although most of the breakdown undoubtedly occurs during purification (despite the purification being carried out at 4°C and with the inclusion of PVPP and protease inhibitors), some of the breakdown may also occur *in vivo*. Breakdown of enzymes *in vivo* has been observed during the purification of starch branching enzyme from potato tubers. Using fresh harvested tubers for the purification resulted in a predominately high molecular weight starch branching enzyme being isolated, but when stored tubers were used, a wide range of molecular weight proteins were isolated (Blennow & Johansson, 1991 *Phytochem.* 30, 437-444).

The fact that the antiserum to SSS recognises the 140 kDa protein in both Estima and Desiree tubers suggests that there is no difference between the two cultivars in their major starch synthases, and vindicates the use of these two different cultivars in the work reported herein. The occurrence of a 100 kDa protein antigenically related to the 140 kDa protein in Desiree tubers is interesting, and at present it is not known what that protein may be. Its absence from Estima tubers could reflect the fact that these cultivars were stored rather than developing tubers, or could represent a difference between cultivars.

Specific activities of the purified soluble starch synthases from potato tuber are comparable with or greater than those of soluble starch synthases from other storage organs. Purification to homogeneity of isoforms of soluble starch synthase resulted in specific activities of 16 μ mol.(mg protein)⁻¹.min⁻¹ from pea embryo (Denyer & Smith 1992 cited previously), 14 μ mol.(mg protein)⁻¹.min⁻¹ from wheat (Denyer *et al.*, 1995 cited previously) and 9 μ mol.(mg protein)⁻¹.min⁻¹ from maize (Mu *et al.* (1994) *Plant J.* 6, 151-159). The specific activity of the soluble starch synthase reported in this application is 7- to 300-fold higher than that of the partial purifications of soluble starch synthase activity from potato tuber reported by Hawker *et al.*, 1972 *Phytochem.* 11, 1278-1293: 0.64 μ mol.(mg protein)⁻¹.min⁻¹, Baba *et al.* (1990) *Phytochem.* 29, 719-723: 0.03 μ mol.(mg protein)⁻¹.min⁻¹ and Ponstein (1.35 and 0.91 μ mol.(mg protein)⁻¹.min⁻¹).

The immunoprecipitation experiments also suggest that the purified proteins are the major soluble starch synthases in potato tuber, or are products directly derived from such synthases. The antiserum raised against the soluble starch

synthase from potato precipitates 75 % of the total soluble starch-synthase activity in crude extract (Figure 5). The remainder of the activity is partly due to GBSS II (Table 2), but the possibility of further minor isoforms cannot be ruled out.

The purified soluble starch synthase is likely to represent a novel class of starch synthase. It is not related to the major soluble starch synthase in pea embryo (GBSS II), which is clearly related to the minor, soluble 92 kDa GBSS II in potato. The soluble starch synthase is only very weakly recognised by the antibody raised to GBSS II from pea. It is not related to the GBSS I proteins either: the starch synthase from potato tuber is not recognised by the antibody raised to GBSS I from pea embryo. These results reinforce the view that storage organs differ profoundly in the nature and number of active isoforms of starch synthase (Smith *et al.*, 1995 Plant Physiol. 107, 1; Edwards *et al.*, and Denyer *et al.*, both cited previously).

Example 2

ISOLATION OF A cDNA CLONE FOR SOLUBLE STARCH SYNTHASE FROM POTATO TUBERS

The antiserum raised to the purified starch synthase protein from Estima tubers was used for immunoscreening of a λ gt 11 library (provided by C Grierson, John Innes Centre, Norwich) containing cDNA inserts with *Eco*RI linkers, constructed from developing Estima tuber poly(A) RNA.

Approximately 1.5×10^6 plaque-forming units were probed with the antiserum at a dilution of 1/1000. The second antibody was an anti-rat immunoglobulin linked to horseradish peroxidase (Amersham International, Amersham, UK). Two positive clones were isolated. These were both 1.1 kb in length and contained poly(A) tracts at their 3' ends. One of these was cloned into the *Eco*RI site of pBluescript SK+ to give plasmid pRAT2. A 5' *Eco*RI-*ECORV* fragment from this clone was used as a probe on the λ gt11 library. Filters were washed in 0.1 x SSC (1 x SSC is 0.15 M NaCl, 0.015 M sodium citrate), 0.5 g L⁻¹ SDS at 65°C. Seven clones of 1.3, 1.53, 1.75, 1.88, 2.15, 2.21, and 2.4 kb were isolated. The longest clone was subcloned as an *Eco*RI fragment into pBluescript SK+ to give plasmid pRAT20. A 600-bp 5' fragment from pRAT20 was used to probe a random primed λ gt11 library prepared from cDNA of developing tubers. Three positive clones were isolated. The longest was 2.3 kb and was subcloned as an *Eco*RI fragment into pBluescript SK+ to give pRAT24.

The 2.3 and 2.4kb partial clones overlapped. The full-length composite cDNA was 4.127kb. The DNA sequence of the full length cDNA, and the predicted polypeptide sequence, are shown in Figure 6. DNA sequences were determined according to Sanger *et al.* (1977) by using Sequenase™ (United States Biochemical). Sequence data were analysed using the Genetics Computer Group (Madison, WI) computer program (Devereux *et al.* 1984 Nucl. Acids Res. 12, 387-395).

To check the identity of the cDNA, the amino acid sequence it predicted was compared with amino acid sequences of two peptides obtained by digestion with endoproteinase Lys-C of the 110-kD protein purified from tubers of cultivar Estima. The peptide sequences FIPIPYTSENVVEGK (Seq. ID No. 1) and HIPVFGG (Seq. ID No. 2) corresponded precisely to predicted sequences from the clone. Attempts to obtain N-terminal amino acid sequence of the purified proteins for comparison with the sequence predicted from the cDNA clone were unsuccessful.

On RNA gel blots of poly(A)+RNA from developing tubers, a partial cDNA clone recognised a single transcript of ~4 kb. This size is considerably greater than those of the transcripts for GBSSI and GBSSII and is consistent with the transcript encoding a protein in the range of 110 to 140 kD.

The deduced amino acid sequence of the soluble starch synthase revealed a protein of 1230 amino acids and a predicted size of 139 kD (Figure 6). At the N terminus was a sequence of ~60 amino acids rich in serine and basic residues and low in acidic residues, which is typical of a chloroplast transit peptide. Based on the consensus of Gavel and von Heinje (1990 FEBS Lett. 261, 455-458), the most likely cleavage site would be between amino acids 60 (Cys) and 61 (Ala), because the serine-rich region ends before this point. Cleavage in this region would give a mature protein of ~132 kD. The structure is somewhat similar to that of GBSSII in that it contains a C-terminal region homologous with starch synthases and bacterial glycogen synthases and an N-terminal extension. The N-terminal extension shows little sequence similarity to the N-terminal extensions of GBSSII from pea or potato (in turn, they show little similarity to each other: Edwards *et al.*, (1995) Plant J. 8, 283-294) or to any other sequence in the data bases. The N-terminal domain resembles those of pea and potato GBSSII in that it shows considerable predicted flexibility (Chou-Fasman algorithm: see Dry *et al.*, (1992) Plant J. 2, 193-202); all these extensions may therefore serve similar roles. At the C-terminal end of the N-terminal extension of the soluble starch synthase are two proline residues; multiple proline residues have been noted previously at the C-terminal ends of N-terminal extensions of both starch synthases and starch-branching enzymes (Dry *et al.*, (1992) Plant J. 2, 193-202; Burton *et al.*, 1995).

The roles of these N-terminal extensions are not known, but it seems likely that they are involved in determining properties such as interaction with starch polymers rather than contributing to basic catalytic properties. The C-terminal region from amino acid 780 to the end shows greatest similarity to glycogen synthases from bacteria, although there

is also similarity to other starch synthases from plants. The KTGG motif close to the N terminus of this region beginning with position 794 is conserved (KVGGL). This domain is thought to be involved in ADP/ADP-glucose binding (Furukawa *et al.*, 1990 J. Biol. Chem. **265**, 2086-2090). Interestingly, a second domain with a similar structure is also conserved in the C termini of all bacterial glycogen synthases and starch synthases (including the motif beginning at position 1143, T/V GGLXDT I/V); this may represent a second domain involved in ADP/ADP-glucose binding. The whole region around this second domain is widely conserved among α -1,4-glucosyltransferases, indicating close involvement with the catalytic process.

Over the rest of the soluble starch synthase protein, there are several other domains showing conservation between different starch synthases. However, it also shows some notable gaps in its sequence when aligned with GBSSI and GBSII, for example, between amino acids 828 to 829 (13 amino acids), 894 to 895 (10 amino acids), and 944 to 945 (35 amino acids). These regions may confer specific properties on GBSSI and GBSII compared with the soluble synthase.

Example 3

POTATO TRANSFORMATION

Binary vectors containing a partial cDNA for soluble starch synthase ("SSS") in the antisense orientation, under the control of a) the double 35S promoter or b) the patatin promoter have been constructed. The 2 x 35S construct is detailed below.

Construction of Antisense Binary Vector

The 1.1-kb *Pst*-*EcoRV* fragment from pRAT2 encoding the 3' end of the soluble starch synthase was subcloned in an antisense orientation between the cauliflower mosaic virus double 35S promoter and cauliflower mosaic virus terminator (*Pst*I-*Sma*I) in pJIT60 (Guerineau and Mullineaux, 1993 "Plant transformation and expression vectors". In Plant Molecular Biology Labfax R.R.D. Croy, Ed. (Oxford, UK BIOS Scientific Publishers) p121-148), producing pRAT3. The *Xho*I-partial *Sst*I fragment from pRAT3, encompassing the promoter, antisense cDNA, and terminator, was ligated between the *Sal*I-*Sst*I sites of the plant transformation vector pBIN19 (Bevan, 1984 Nucl. Acids Res. **12**, 8711-8721), resulting in plasmid pRAT4. This plasmid is illustrated schematically in Figure 7, which Figure also shows the plasmid pPATRAT comprising the patatin promoter.

Transformation of Potato

Binary plasmid pRAT4 was introduced into *Agrobacterium tumefaciens* by the freeze-thaw method of An *et al.*, (1988 Binary Vectors. In Plant Molecular Biology Manual A3. Eds. Gelvin S.B. and Schilperoort R.A. ppl-19). Preparation of *Agrobacterium* inoculum carrying the antisense construct, inoculation of tuber discs of potato cultivar Desiree, regeneration of shoots, and rooting of shoots were as described by Edwards *et al.* (1995 Plant J. **8**, 283-294).

Thirteen independently transformed plants and four independent control plants (transformed with the vector alone) were transferred to a soil-based compost and allowed to develop tubers. The presence of the SSS antisense construct was confirmed by DNA gel blotting (data not shown). Six of the transgenic plants had levels of SSS transcript indistinguishable from those of the control plants on RNA gel blots. However, seven independent transformants (named 1, 2, 9, 18, 19, 25 and 26) had strongly reduced or undetectable levels of SSS transcript. The loss or reduction of detectable transcript was specific for SSS, and there was little variation in the level of transcript for GBSSI among the plants studied (data shown in Marshall *et al.*, 1996 The Plant Cell **8**, 1121-1135).

Tubers of the transformants with unaltered levels of SSS transcript had soluble starch synthase activities that were indistinguishable from those of the control plants and from values typical of those obtained from developing Desiree tubers in general (Edwards *et al.* (1995) Plant J. **8**, 283-294 1995). Tubers of the seven transformants with reduced or undetectable levels of SSS transcript had significantly reduced activities, and three plants displayed activities that were 30% or less of the average value of the control plants. Table 3 shows that the observed reductions in soluble starch synthase activity were reproducible from one tuber to another. They were also reproducible through tuber development.

Table 3. Effects of Reduced Activity of SSS on Soluble and Granule-Bound Starch Synthase Activity and Amylose Content of Starch.

Plant ^a	Soluble Activity ^b (nmol min ⁻¹ g ⁻¹ Fresh Weight)	Granule-Bound Activity ^c (nmol min ⁻¹ g ⁻¹ Fresh Weight)	Amylose Content ^d (% Total Starch)
1	ND ^e	ND	27.8
2	ND	ND	29.5, 29.8
9	18.3 \pm 3.9 (4)	118	28.6
18	23.6 \pm 6.7 (3)	97	29.3
25	29.5 \pm 3.6 (4)	113	27.3
26	33.3 \pm 8.3 (3)	80	30.1
Control	98.4 \pm 4.9 (9)	106 \pm 12	26.4, 28.9
Desiree	ND	ND	27.8, 29.2

^a Plant numbers refer to individual transgenic plants with reduced SSS activity. Tubers are from a single plant, except for the control line, in which three different plants (each an independent, control transformant) were used.

^b Soluble activity was measured by using duplicate samples from tubers of 12 to 70 g fresh weight harvested at intervals during plant development. Values are the means \pm SE of measurements made with the number of tubers given within parentheses.

^c Granule-bound activities are the means of measurements made by using duplicate samples from a single tuber (12 to 70 g fresh weight) harvested at maturity.

^d Amylose content was measured by using starch extracted from two or three tubers per mature plant. Values are the means of measurements made with two separate samples taken from the bulk starch preparations: two values are given when independent starch preparations were used. Wild-type Desiree plants used for these measurements were grown in the same greenhouse at the same time as the transgenic

plants.

^c ND, not determined.

Reductions in Starch Synthase Activity Are Specifically Due to Loss of SSS

To discover whether the reductions in activity were specifically attributable to loss of SSS, two sorts of experiments were undertaken. First, isoforms were visualised on native gels of crude, soluble extracts of transformed tubers. The group of bands attributable to SSS was present in extracts from control plants and from all six of the transformants with soluble starch synthase activities comparable with control activities. It was absent from extracts of all seven transformants with reduced starch synthase activities. Other groups of bands on the gels, including those attributable to GBSSII, were present in all extracts (data shown in Marshall *et al.*, 1996).

Second, crude, soluble extracts from a plant with strongly reduced activity were incubated with the antiserum raised against SSS. The antiserum inhibited activity by 16%, compared with 75% inhibition in extracts of untransformed tubers of cultivar Désirée (Table 2).

Loss of starch synthase activity from the soluble fraction in transgenic tubers was accompanied by dramatic reductions in the amount of the 140-kD protein recognised by the antiserum in soluble and granule-bound fractions of the tuber. The protein was not detected, or detected only very weakly relative to controls, on immunoblots of these fractions from tubers of the six transgenic lines with the largest reductions in starch synthase activity. In contrast, the soluble protein of 105 kD also recognised by the antiserum was present in equal amounts in all lines examined (data shown in Marshall *et al.*, 1996).

Reduction in SSS Activity Alters Granule Shape but Has Little Effect on Starch and Amylose Content

Tubers of the seven transformants with reduced activities of soluble starch synthase had starch granules with strikingly altered morphology. Two types of granule were present: simple granules with deep, often T-shaped cracks centered on the hilum, and granules that appeared to be large clusters of tiny, spherical granules. A range of different sizes of both types of granule was present in tubers at various developmental stages (data not shown).

In spite of the alteration in granule morphology, tubers of transformants with reduced activity of SSS were indistinguishable from control tubers with respect to total starch content. This was true of both developing tubers and tubers of mature plants on which the haulm was senescent. The starch of these plants also displayed no significant alteration in amylose content (Table 3).

Reduction in SSS Activity Does Not Affect Other Isoforms of Starch Synthase

It was thought possible that the reduction in SSS in transformed tubers may have secondary effects on other isoforms of starch synthase. Any alterations in other isoforms could seriously affect deductions about the importance and role of SSS and might prevent alteration of starch properties in transformed plants. Effects of the reduction in SSS on GBSSI were assessed by measuring granule-bound starch synthase activity in crude extracts of tubers and examining gels of granule-bound proteins. There was no difference in granule-bound activity between control plants and those in which soluble starch synthase activity was reduced (Table 3). More than 95% of the starch synthase activity of intact starch granules of wild-type potatoes is attributable to GBSSI (Edwards *et al.* (1995) Plant J. 8, 283-294 1995). Reductions in SSS also had no obvious effect on the amount of GBSSI protein bound to starch granules (data not shown).

Effects of reductions in SSS on GBSSII were assessed in three ways. First, amounts of GBSSII protein in the soluble and granule-bound fractions of the tuber were visualised by immunoblotting. There were no obvious differences between control plants and those in which SSS was reduced.

Second, as described above, GBSSII was visualised on native gels of crude, soluble extracts stained for starch synthase activity. Again, there were no marked or consistent differences between control plants and those in which SSS was reduced.

Third, immunoprecipitation experiments were used to assess the proportion of the residual activity attributable to GBSSII in tubers in which SSS was reduced. The antiserum raised against GBSSII of pea embryos, which recognises GBSSII of potatoes (Edwards *et al.* 1995 cited above), inhibited ~40% of the activity in tubers in which soluble starch synthase activity was reduced by ~80% (line 9) compared with 9% in control and wild-type tubers (Table 2). Using these figures and starch synthase activities from Table 3, the activity attributable to GBSSII is 7.3 nmol min⁻¹g⁻¹ fresh

weight in line 9 and $8.8 \text{ nmol min}^{-1} \text{g}^{-1}$ fresh weight in control tubers. This indicates that the reduction in SSS has little effect on the soluble activity of GBSSII.

Example 4

5

Detailed analysis of starch from tubers obtained from transformed Potato plants

Despite the results of crude analysis described in Example 3, indicating that the starch from transformed plants was essentially unaltered, it was decided to perform more detailed analysis of the starch, by Differential Scanning Calorimetry and Viscoamylograph. Analysis was performed as described in WO 95/26407 and WO 96/34968.

Surprisingly it was found that certain physical properties of the starch were consistently significantly altered. In particular, it was found that the viscosity onset temperature was significantly reduced compared to starch obtained from equivalent control plants which did not contain the SSS antisense construct. The results are shown in Table 4.

15

20

25

30

35

40

45

50

55

Starch from AS major soluble starch synthase tubers

TABLE 4

Plant line	Plant n°	DSC		RVA	SSS activity (nmol/min/ mg of tuber)	Apparent amylose* (% w/w)	Granule morphology Comments
		Peak (°C)	Onset (°C)	Delta H (J/g)			
Rat 4.1		65.8	61.7	16.1		27.82	unusual granules: some compound, some with large cracks
Rat 4.2	1	65.5	61.6	16.0		29.76	unusual granules (as Rat 4.1)
	2	63.4	59.4	17.0		29.45	DSC endotherm double peak
		63.6	59.6	16.7			
Desiree (control)	1	64.0	60.1	16.9		29.16	
	2	63.1	59.3	17.4		27.80	
		67.6	64.4	17.5			
		67.5	64.4	17.2			
		67.3	64.6	10.9			
		67.3	64.7	17.5			
Rat 4.9		57.4	53.2	14.5	18.35	28.60	unusual granules (as Rat 4.1)
Rat 4.18		57.4	53.2	14.0		29.31	unusual granules (as Rat 4.1), but smaller than controls. DSC endotherm broad
Rat 4.25		58.6	54.3	(14.5)	23	27.32	unusual granules (as Rat 4.1)
		58.5	54.3	(16.0)	24		
		59.0	54.3	14.8			
		59.0	54.2	14.9			
Rat 4.26		(59.6)	54.9	(16.2)	30	30.10	unusual granules (as Rat 4.1)
		(59.8)	54.9	(16.9)			
Luc 1 (control)		63.7	61.0	17.1	110	26.43	DSC endotherm double peak
Luc 6 (control)		63.6	61.3	17.6			
		64.6	62.0	17.9	116	28.85	
		64.5	62.0	17.3			

DSC Differential scanning calorimetry

Onset: endotherm onset temperature
 Peak: endotherm peak temperature
 Delta H: endotherm enthalpy

RVA

RVA Rapid visco analyser
 Onset: viscosity onset temperature

SSS activity

Soluble starch synthase activity

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

(A) NAME: National Starch and Chemical Investment
Holding Corporation
(B) STREET: Suite 27, 501 Silverside Road
(C) CITY: Wilmington
(D) STATE: Delaware
(E) COUNTRY: U.S.A.
(F) POSTAL CODE (ZIP): DE 19809

(ii) TITLE OF INVENTION: Improvements in or Relating to Soluble
Starch Synthase

(iii) NUMBER OF SEQUENCES: 4

(iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

Phe	Ile	Pro	Ile	Pro	Tyr	Thr	Ser	Glu	Asn	Val	Val	Glu	Gly	Lys
1				5					10					15

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 7 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

His Ile Pro Val Phe Gly Gly
1 5

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 4127 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Solanum tuberosum
(B) STRAIN: Desiree
(F) TISSUE TYPE: tuber

(vii) IMMEDIATE SOURCE:

(A) LIBRARY: lambda gt11

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION:143..3835

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
(B) LOCATION:143..322

(ix) FEATURE:

(A) NAME/KEY: mat_peptide
(B) LOCATION:323..3832

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

GAATTCGCGG CCGCAGATAG TGTGTTGATG AAGGAGAAGA GAGATATTTT ACATGGGATG 60

TTCTATTTGA TTCTGTGGTG AACAAAGATT TTACAAAGAA CATTCCTTTT TCTTTTTTCC 120

TTGGTTCTTG TGTGGGTCAG CC ATG GAT GTT CCA TTT CCA CTG CAT AGA TCA 172

Met Asp Val Pro Phe Pro Leu His Arg Ser
-60 -55

TTG AGT TGC ACA AGT GTC TCC AAT GCA ATA ACC CAC CTC AAG ATC AAA 220

Leu Ser Cys Thr Ser Val Ser Asn Ala Ile Thr His Leu Lys Ile Lys
-50 -45 -40 -35

	CCT ATT CTT GGG TTT GTC TCT CAT GGA ACC ACA AGT CTA TCA GTA CAA	268
	Pro Ile Leu Gly Phe Val Ser His Gly Thr Thr Ser Leu Ser Val Gln	
	-30 -25 -20	
5	TCT TCT TCA TGG AGG AAG GAT GGA ATG GTT ACT GGG GTT TCA TTT TCC	316
	Ser Ser Ser Trp Arg Lys Asp Gly Met Val Thr Gly Val Ser Phe Ser	
	-15 -10 -5	
10	ATT TGT GCA AAT TTC TCG GGA AGA AGA CGG AGA AAA GTT TCA ACT CCT	364
	Ile Cys Ala Asn Phe Ser Gly Arg Arg Arg Arg Lys Val Ser Thr Pro	
	1 5 10	
15	AGG AGT CAA GGC TCT TCA CCT AAG GGG TTT GTG CCA AGG AAG CCC TCA	412
	Arg Ser Gln Gly Ser Ser Pro Lys Gly Phe Val Pro Arg Lys Pro Ser	
	15 20 25 30	
20	GGG ATG AGC ACG CAA AGA AAG GTT CAG AAG AGC AAT GGT GAT AAA GAA	460
	Gly Met Ser Thr Gln Arg Lys Val Gln Lys Ser Asn Gly Asp Lys Glu	
	35 40 45	
25	AGT AAA AGT ACT TCA ACA TCT AAA GAA TCT GAA ATT TCC AAC CAG AAG	508
	Ser Lys Ser Thr Ser Thr Ser Lys Glu Ser Glu Ile Ser Asn Gln Lys	
	50 55 60	
30	ACG GTT GAA GCA AGA GTT GAA ACT AGT GAC GAT GAC ACT AAA GGA GTG	556
	Thr Val Glu Ala Arg Val Glu Thr Ser Asp Asp Asp Thr Lys Gly Val	
	65 70 75	
35	GTG AGG GAC CAC AAG TTT CTG GAG GAT GAG GAT GAA ATC AAT GGT TCT	604
	Val Arg Asp His Lys Phe Leu Glu Asp Glu Asp Glu Ile Asn Gly Ser	
	80 85 90	
40	ACT AAA TCA ATA AGT ATG TCA CCT GTT CGT GTA TCA TCT CAA TTT GTT	652
	Thr Lys Ser Ile Ser Met Ser Pro Val Arg Val Ser Ser Gln Phe Val	
	95 100 105 110	
45	GAA AGT GAA GAA ACT GGT GGT GAT GAC AAG GAT GCT GTA AAG TTA AAC	700
	Glu Ser Glu Glu Thr Gly Gly Asp Asp Lys Asp Ala Val Lys Leu Asn	
	115 120 125	
50	AAA TCA AAG AGA TCG GAA GAG AGT GGT TTT ATA ATT GAT TCT GTA ATA	748
	Lys Ser Lys Arg Ser Glu Glu Ser Gly Phe Ile Ile Asp Ser Val Ile	
	130 135 140	
55	AGA GAA CAA AGT GGA TCT CAG GGG GAA ACT AAT GCC AGT AGC AAG GGA	796
	Arg Glu Gln Ser Gly Ser Gln Gly Glu Thr Asn Ala Ser Ser Lys Gly	
	145 150 155	
60	AGC CAT GCT GTG GGT ACA AAA CTT TAT GAG ATA TTG CAG GTG GAT GTT	844
	Ser His Ala Val Gly Thr Lys Leu Tyr Glu Ile Leu Gln Val Asp Val	
	160 165 170	
65	GAG CCA CAA CAA TTG AAA GAA AAT AAT GCT GGG AAT GTT GAA TAC AAA	892
	Glu Pro Gln Gln Leu Lys Glu Asn Asn Ala Gly Asn Val Glu Tyr Lys	
	175 180 185 190	

	GGA CCT GAA GCA AGT AAG CTA TTG GAA ATT ACT AAG GCT AGT GAT GTG	940
	Gly Pro Val Ala Ser Lys Leu Leu Glu Ile Thr Lys Ala Ser Asp Val	
	195 200 205	
5	GAA CAC ACT GAA AGC AAT GAG ATT GAT GAC TTA GAC ACT AAT AGT TTC	988
	Glu His Thr Glu Ser Asn Glu Ile Asp Asn Leu Asp Thr Asn Ser Phe	
	210 215 220	
10	TTT AAA TCA GAT TTA ATT GAA GAG GAT GAG CCA TTA GCT GCA GGA ACA	1036
	Phe Lys Ser Asp Leu Ile Glu Glu Asp Glu Pro Leu Ala Ala Gly Thr	
	225 230 235	
15	GTG GAG ACT GGA GAT TCT TCT CTA AAC TTA AGA TTG GAG ATG GAA GCA	1084
	Val Glu Thr Gly Asp Ser Ser Leu Asn Leu Arg Leu Glu Met Glu Ala	
	240 245 250	
20	AAT CTA CGT AGG CAG GCT ATA GAA AGG CTT GCC GAG GAA AAT TTA TTG	1132
	Asn Leu Arg Arg Gln Ala Ile Glu Arg Leu Ala Glu Glu Asn Leu Leu	
	255 260 265 270	
	CAA GGG ATC AGA TTA TTT TGT TTT CCA GAG GTT GTA AAA CCT GAT GAA	1180
	Gln Gly Ile Arg Leu Phe Cys Phe Pro Glu Val Val Lys Pro Asp Glu	
	275 280 285	
25	GAT GTC GAG ATA TTT CTT AAC AGA GGT CTT TCC ACT TTG AAG AAT GAG	1228
	Asp Val Glu Ile Phe Leu Asn Arg Gly Leu Ser Thr Leu Lys Asn Glu	
	290 295 300	
30	TCT GAT GTC TTG ATT ATG GGA GCT TTT AAT GAG TGG CGC TAT AGG TCT	1276
	Ser Asp Val Leu Ile Met Gly Ala Phe Asn Glu Trp Arg Tyr Arg Ser	
	305 310 315	
35	TTT ACT ACA AGG CTA ACT GAG ACT CAT CTC AAT GGA GAT TGG TGG TCT	1324
	Phe Thr Thr Arg Leu Thr Glu Thr His Leu Asn Gly Asp Trp Trp Ser	
	320 325 330	
40	TGC AAG ATC CAT GTT CCC AAG GAA GCA TAC AGG GCT GAT TTT GTG TTT	1372
	Cys Lys Ile His Val Pro Lys Glu Ala Tyr Arg Ala Asp Phe Val Phe	
	335 340 345 350	
	TTT AAT GGA CAA GAT GTC TAT GAC AAC AAT GAT GGA AAT GAC TTC AGT	1420
	Phe Asn Gly Gln Asp Val Tyr Asp Asn Asn Asp Gly Asn Asp Phe Ser	
	355 360 365	
45	ATA ACT GTG AAA GGT GGT ATG CAA ATC ATT GAC TTT GAA AAT TTC TTG	1468
	Ile Thr Val Lys Gly Gly Met Gln Ile Ile Asp Phe Glu Asn Phe Leu	
	370 375 380	
50	CTT GAG GAG AAA TGG AGA GAA CAG GAG AAA CTT GCT AAA GAA CAA GCT	1516
	Leu Glu Glu Lys Trp Arg Glu Gln Glu Lys Leu Ala Lys Glu Gln Ala	
	385 390 395	
55	GAA AGA GAA AGA CTA GCA GAA GAA CAA AGA CGA ATA GAA GCA GAG AAA	1564
	Glu Arg Glu Arg Leu Ala Glu Glu Gln Arg Arg Ile Glu Ala Glu Lys	
	400 405 410	

5	GCT GAA ATT GAA GCT GAC AGA GCA CAA GCA AAG GAA GAG GCT GCA AAG 1612 Ala Glu Ile Glu Ala Asp Arg Ala Gln Ala Lys Glu Glu Ala Ala Lys 415 420 425 430
10	AAA AAG AAA GTA TTG CGA GAA TTG ATG GTA AAA GCC ACG AAG ACT CGT 1660 Lys Lys Lys Val Leu Arg Glu Leu Met Val Lys Ala Thr Lys Thr Arg 435 440 445
15	GAT ATC ACC TGG TAC ATA GAG CCA AGT GAA TTT AAA TGC GAG GAC AAG 1708 Asp Ile Thr Trp Tyr Ile Glu Pro Ser Glu Phe Lys Cys Glu Asp Lys 450 455 460
20	GTC AGG TTA TAC TAT AAC AAA AGT TCA GGT CCT CTC TCC CAT GCT AAG 1756 Val Arg Leu Tyr Tyr Asn Lys Ser Ser Gly Pro Leu Ser His Ala Lys 465 470 475
25	GAC TTG TGG ATC CAC GGA GGA TAT AAT AAT TGG AAG GAT GGT TTG TCT 1804 Asp Leu Trp Ile His Gly Gly Tyr Asn Asn Trp Lys Asp Gly Leu Ser 480 485 490
30	ATT GTC AAA AAG CTT GTT AAA TCT GAG AGA ATA GAT GGT GAT TGG TGG 1852 Ile Val Lys Lys Leu Val Lys Ser Glu Arg Ile Asp Gly Asp Trp Trp 495 500 505 510
35	TAT ACA GAG GTT GTT ATT CCT GAT CAG GCA CTT TTC TTG GAT TGG GTT 1900 Tyr Thr Glu Val Val Ile Pro Asp Gln Ala Leu Phe Leu Asp Trp Val 515 520 525
40	TTT GCT GAT GGT CCA CCC AAG CAT GCC ATT GCT TAT GAT AAC AAT CAC 1948 Phe Ala Asp Gly Pro Pro Lys His Ala Ile Ala Tyr Asp Asn Asn His 530 535 540
45	CGC CAA GAC TTC CAT GCC ATT GTC CCC AAC CAC ATT CCG GAG GAA TTA 1996 Arg Gln Asp Phe His Ala Ile Val Pro Asn His Ile Pro Glu Glu Leu 545 550 555
50	TAT TGG GTT GAG GAA GAA CAT CAG ATC TTT AAG ACA CTT CAG GAG GAG 2044 Tyr Trp Val Glu Glu Glu His Gln Ile Phe Lys Thr Leu Gln Glu Glu 560 565 570
55	AGA AGG CTT AGA GAA GCG GCT ATG CGT GCT AAG GTT GAA AAA ACA GCA 2092 Arg Arg Leu Arg Glu Ala Ala Met Arg Ala Lys Val Glu Lys Thr Ala 575 580 585 590
60	CTT CTG AAA ACT GAA ACA AAG GAA AGA ACT ATG AAA TCA TTT TTA CTG 2140 Leu Leu Lys Thr Glu Thr Lys Glu Arg Thr Met Lys Ser Phe Leu Leu 595 600 605
65	TCT CAG AAG CAT GTA GTA TAT ACT GAA CCT CTT GAT ATC CAA GCT GGA 2188 Ser Gln Lys His Val Val Tyr Thr Glu Pro Leu Asp Ile Gln Ala Gly 610 615 620
70	AGC AGC GTC ACA GTT TAC TAT AAT CCC GCC AAT ACA GTA CTT AAT GGT 2236 Ser Ser Val Thr Val Tyr Tyr Asn Pro Ala Asn Thr Val Leu Asn Gly 625 630 635

5	AAA CCT GAA ATT TGG TTC AGA TGT TCA TTT AAT CGC TGG ACT CAC CGC 2284 Lys Pro Glu Ile Trp Phe Arg Cys Ser Phe Asn Arg Trp Thr His Arg 640 645 650
10	CTG GGT CCA TTG CCA CCT CAG AAA ATG TCG CCT GCT GAA AAT GGC ACC 2332 Leu Gly Pro Leu Pro Pro Gln Lys Met Ser Pro Ala Glu Asn Gly Thr 655 660 665 670
15	CAT GTC AGA GCA ACT GTG AAG GTT CCA TTG GAT GCA TAT ATG ATG GAT 2380 His Val Arg Ala Thr Val Lys Val Pro Leu Asp Ala Tyr Met Met Asp 675 680 685
20	TTT GTA TTT TCC GAG AGA GAA GAT GGT GGG ATT TTT GAC AAT AAG AGC 2428 Phe Val Phe Ser Glu Arg Glu Asp Gly Gly Ile Phe Asp Asn Lys Ser 690 695 700
25	GGA ATG GAC TAT CAC ATA CCT GTG TTT GGA GGA GTC GCT AAA GAA CCT 2476 Gly Met Asp Tyr His Ile Pro Val Phe Gly Gly Val Ala Lys Glu Pro 705 710 715
30	CCA ATG CAT ATT GTC CAT ATT GCT GTC GAA ATG GCA CCA ATT GCA AAG 2524 Pro Met His Ile Val His Ile Ala Val Glu Met Ala Pro Ile Ala Lys 720 725 730
35	GTG GGA GGC CTT GGT GAT GTT GTT ACT AGT CTT TCC CGT GCT GTT CAA 2572 Val Gly Gly Leu Gly Asp Val Val Thr Ser Leu Ser Arg Ala Val Gln 735 740 745 750
40	GAT TTA AAC CAT AAT GTG GAT ATT ATC TTA CCT AAG TAT GAC TGT TTG 2620 Asp Leu Asn His Asn Val Asp Ile Ile Leu Pro Lys Tyr Asp Cys Leu 755 760 765
45	AAG ATG AAT AAT GTG AAG GAC TTT CGG TTT CAC AAA AAC TAC TTT TGG 2668 Lys Met Asn Asn Val Lys Asp Phe Arg Phe His Lys Asn Tyr Phe Trp 770 775 780
50	GGT GGG ACT GAA ATA AAA GTA TGG TTT GGA AAG GTG GAA GGT CTC TCG 2716 Gly Gly Thr Glu Ile Lys Val Trp Phe Gly Lys Val Glu Gly Leu Ser 785 790 795
55	GTC TAT TTT TTG GAG CCT CAA AAC GGG TTA TTT TCG AAA GGG TGC GTC 2764 Val Tyr Phe Leu Glu Pro Gln Asn Gly Leu Phe Ser Lys Gly Cys Val 800 805 810
60	TAT GGT TGT AGC AAC GAT GGT GAA CGA TTT GGT TTC TTC TGT CAC GCG 2812 Tyr Gly Cys Ser Asn Asp Gly Glu Arg Phe Gly Phe Phe Cys His Ala 815 820 825 830
65	GCT TTG GAG TTT CTT CTG CAA GGT GGA TTT AGT CCG GAT ATC ATT CAT 2860 Ala Leu Glu Phe Leu Leu Gln Gly Gly Phe Ser Pro Asp Ile Ile His 835 840 845
70	TGC CAT GAT TGG TCT AGT GCT CCT GTT GCT TGG CTC TTT AAG GAA CAA 2908 Cys His Asp Trp Ser Ser Ala Pro Val Ala Trp Leu Phe Lys Glu Gln 850 855 860

TAT ACA CAC TAT GGT CTA AGC AAA TCT CGT ATA GTC TTC ACG ATA CAT 2956
 Tyr Thr His Tyr Gly Leu Ser Lys Ser Arg Ile Val Phe Thr Ile His
 365 870 875

5 AAT CTT GAA TTT GGG GCA GAT CTC ATT GGG AGA GCA ATG ACT AAC GCA 3004
 Asn Leu Glu Phe Gly Ala Asp Leu Ile Gly Arg Ala Met Thr Asn Ala
 880 885 890

10 GAC AAA GCT ACA ACA GTT TCA CCA ACT TAC TCA CAG GAG GTG TCT GGA 3052
 Asp Lys Ala Thr Thr Val Ser Pro Thr Tyr Ser Gln Glu Val Ser Gly
 895 900 905 910

15 AAC CCT GTA ATT GCG CCT CAC CTT CAC AAG TTC CAT GGT ATA GTG AAT 3100
 Asn Pro Val Ile Ala Pro His Leu His Lys Phe His Gly Ile Val Asn
 915 920 925

20 GGG ATT GAC CCA GAT ATT TGG GAT CCT TTA AAC GAT AAG TTC ATT CCG 3148
 Gly Ile Asp Pro Asp Ile Trp Asp Pro Leu Asn Asp Lys Phe Ile Pro
 930 935 940

25 ATT CCG TAC ACT TCA GAA AAC GTT GTT GAG GGC AAA ACA GCA GCC AAG 3196
 Ile Pro Tyr Thr Ser Glu Asn Val Val Glu Gly Lys Thr Ala Ala Lys
 945 950 955

30 GAA GCT TTG CAG CGA AAA CTT GGA CTG AAA CAG GCT GAC CTT CCT TTG 3244
 Glu Ala Leu Gln Arg Lys Leu Gly Leu Lys Gln Ala Asp Leu Pro Leu
 960 965 970

35 GTA GGA ATT ATC ACC CGC TTA ACT CAC CAG AAA GGA ATC CAC CTC ATT 3292
 Val Gly Ile Ile Thr Arg Leu Thr His Gln Lys Gly Ile His Leu Ile
 975 980 985 990

40 AAA CAT GCT ATT TGG CGC ACC TTG GAA CGG AAC GGA CAG GTA GTC TTG 3340
 Lys His Ala Ile Trp Arg Thr Leu Glu Arg Asn Gly Gln Val Val Leu
 995 1000 1005

45 CTT GGT TCT GCT CCT GAT CCT AGG GTA CAA AAC AAT TTT GTT AAT TTG 3388
 Leu Gly Ser Ala Pro Asp Pro Arg Val Gln Asn Asn Phe Val Asn Leu
 1010 1015 1020

50 GCA AAT CAA TTG CAC TCC AAA TAT AAT GAC CGC GCA CGA CTC TGT CTA 3436
 Ala Asn Gln Leu His Ser Lys Tyr Asn Asp Arg Ala Arg Leu Cys Leu
 1025 1030 1035

55 ACA TAT GAC GAG CCA CTT TCT CAC CTG ATA TAT GCT GGT GCT GAT TTT 3484
 Thr Tyr Asp Glu Pro Leu Ser His Leu Ile Tyr Ala Gly Ala Asp Phe
 1040 1045 1050

60 ATT CTA GTT CCT TCA ATA TTT GAG CCA TGT GGA CTA ACA CAA CTT ACC 3532
 Ile Leu Val Pro Ser Ile Phe Glu Pro Cys Gly Leu Thr Gln Leu Thr
 1055 1060 1065 1070

65 GCT ATG AGA TAT GGT TCA ATT CCA GTC GTG CGT AAA ACT GGA GGA CTT 3580
 Ala Met Arg Tyr Gly Ser Ile Pro Val Val Arg Lys Thr Gly Gly Leu
 1075 1080 1085

TAT GAT ACT GTA TTT GAT GTT GAC CAT GAC AAA GAG AGA GCA CAA CAG 3628
 Tyr Asp Thr Val Phe Asp Val Asp His Asp Lys Glu Arg Ala Gln Gln
 1090 1095 1100
 5
 TGT GGT CTT GAA CCA AAT GGA TTC AGC TTT GAT GGA GCA GAT GCT GGC 3676
 Cys Gly Leu Glu Pro Asn Gly Phe Ser Phe Asp Gly Ala Asp Ala Gly
 1105 1110 1115
 10
 GGA GTT GAT TAT GCT CTG AAT AGA GCT CTC TCT GCT TGG TAC GAT GGT 3724
 Gly Val Asp Tyr Ala Leu Asn Arg Ala Leu Ser Ala Trp Tyr Asp Gly
 1120 1125 1130
 15
 CGG GAT TGG TTC AAC TCT TTA TGC AAG CAG GTC ATG GAA CAA GAT TGG 3772
 Arg Asp Trp Phe Asn Ser Leu Cys Lys Gln Val Met Glu Gln Asp Trp
 1135 1140 1145 1150
 20
 TCT TGG AAC CGA CCT GCT CTT GAT TAT TTG GAG CTT TAC CAT GCT GCT 3820
 Ser Trp Asn Arg Pro Ala Leu Asp Tyr Leu Glu Leu Tyr His Ala Ala
 1155 1160 1165
 AGA AAG TTA GAA TAG TTAGTTTGTG AGATGCTAGC AGAAAAATTC ACGAGATCTG 3875
 Arg Lys Leu Glu *
 1170
 25
 CAATCTGTAC AGGTTTCAGTG TTGGGTCTG GACAGCTTTA TCATTTCTTA TATCAAAGTA 3935
 TAAATCAAGT CTACACTGAG GATCAATAGC AGACAGTCCT CAAGTTCATT TCATTTTTTTG 3995
 30
 GGGCAAACAT ATGAAAGAGC TTAGCCTCTT AATAATGTCG GCCTATTGAT GATTATTTGT 4055
 TTTGGGAAGA AATGAGAAAT CAAAGGATGC AAAATAAAAA AAAAAAAAAA AAAAAAACT 4115
 CGTGCCGAAT TC 4127

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1231 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Met Asp Val Pro Phe Pro Leu His Arg Ser Leu Ser Cys Thr Ser Val
 -60 -55 -50 -45
 Ser Asn Ala Ile Thr His Leu Lys Ile Lys Pro Ile Leu Gly Phe Val
 -40 -35 -30
 Ser His Gly Thr Thr Ser Leu Ser Val Gln Ser Ser Ser Trp Arg Lys
 -25 -20 -15
 55
 Asp Gly Met Val Thr Gly Val Ser Phe Ser Ile Cys Ala Asn Phe Ser
 -10 -5 1

Gly Arg Arg Arg Arg Lys Val Ser Thr Pro Arg Ser Gln Gly Ser Ser
 5 10 15 20
 5 Pro Lys Gly Phe Val Pro Arg Lys Pro Ser Gly Met Ser Thr Gln Arg
 25 30 35
 Lys Val Gln Lys Ser Asn Gly Asp Lys Glu Ser Lys Ser Thr Ser Thr
 40 45 50
 10 Ser Lys Glu Ser Glu Ile Ser Asn Gln Lys Thr Val Glu Ala Arg Val
 55 60 65
 Glu Thr Ser Asp Asp Asp Thr Lys Gly Val Val Arg Asp His Lys Phe
 70 75 80
 15 Leu Glu Asp Glu Asp Glu Ile Asn Gly Ser Thr Lys Ser Ile Ser Met
 85 90 95 100
 20 Ser Pro Val Arg Val Ser Ser Gln Phe Val Glu Ser Glu Glu Thr Gly
 105 110 115
 Gly Asp Asp Lys Asp Ala Val Lys Leu Asn Lys Ser Lys Arg Ser Glu
 120 125 130
 25 Glu Ser Gly Phe Ile Ile Asp Ser Val Ile Arg Glu Gln Ser Gly Ser
 135 140 145
 Gln Gly Glu Thr Asn Ala Ser Ser Lys Gly Ser His Ala Val Gly Thr
 150 155 160
 30 Lys Leu Tyr Glu Ile Leu Gln Val Asp Val Glu Pro Gln Gln Leu Lys
 165 170 175 180
 Glu Asn Asn Ala Gly Asn Val Glu Tyr Lys Gly Pro Val Ala Ser Lys
 185 190 195
 35 Leu Leu Glu Ile Thr Lys Ala Ser Asp Val Glu His Thr Glu Ser Asn
 200 205 210
 40 Glu Ile Asp Asp Leu Asp Thr Asn Ser Phe Phe Lys Ser Asp Leu Ile
 215 220 225
 Glu Glu Asp Glu Pro Leu Ala Ala Gly Thr Val Glu Thr Gly Asp Ser
 230 235 240
 45 Ser Leu Asn Leu Arg Leu Glu Met Glu Ala Asn Leu Arg Arg Gln Ala
 245 250 255 260
 Ile Glu Arg Leu Ala Glu Glu Asn Leu Leu Gln Gly Ile Arg Leu Phe
 265 270 275
 50 Cys Phe Pro Glu Val Val Lys Pro Asp Glu Asp Val Glu Ile Phe Leu
 280 285 290
 55 Asn Arg Gly Leu Ser Thr Leu Lys Asn Glu Ser Asp Val Leu Ile Met
 295 300 305

Gly Ala Phe Asn Glu Trp Arg Tyr Arg Ser Phe Thr Thr Arg Leu Thr
 310 315 320
 5 Glu Thr His Leu Asn Gly Asp Trp Trp Ser Cys Lys Ile His Val Pro
 325 330 335 340
 Lys Glu Ala Tyr Arg Ala Asp Phe Val Phe Phe Asn Gly Gln Asp Val
 345 350 355
 10 Tyr Asp Asn Asn Asp Gly Asn Asp Phe Ser Ile Thr Val Lys Gly Gly
 360 365 370
 Met Gln Ile Ile Asp Phe Glu Asn Phe Leu Leu Glu Glu Lys Trp Arg
 375 380 385
 15 Glu Gln Glu Lys Leu Ala Lys Glu Gln Ala Glu Arg Glu Arg Leu Ala
 390 395 400
 20 Glu Glu Gln Arg Arg Ile Glu Ala Glu Lys Ala Glu Ile Glu Ala Asp
 405 410 415 420
 Arg Ala Gln Ala Lys Glu Glu Ala Ala Lys Lys Lys Lys Val Leu Arg
 425 430 435
 25 Glu Leu Met Val Lys Ala Thr Lys Thr Arg Asp Ile Thr Trp Tyr Ile
 440 445 450
 Glu Pro Ser Glu Phe Lys Cys Glu Asp Lys Val Arg Leu Tyr Tyr Asn
 455 460 465
 30 Lys Ser Ser Gly Pro Leu Ser His Ala Lys Asp Leu Trp Ile His Gly
 470 475 480
 35 Gly Tyr Asn Asn Trp Lys Asp Gly Leu Ser Ile Val Lys Lys Leu Val
 485 490 495 500
 Lys Ser Glu Arg Ile Asp Gly Asp Trp Trp Tyr Thr Glu Val Val Ile
 505 510 515
 40 Pro Asp Gln Ala Leu Phe Leu Asp Trp Val Phe Ala Asp Gly Pro Pro
 520 525 530
 Lys His Ala Ile Ala Tyr Asp Asn Asn His Arg Gln Asp Phe His Ala
 535 540 545
 45 Ile Val Pro Asn His Ile Pro Glu Glu Leu Tyr Trp Val Glu Glu Glu
 550 555 560
 50 His Gln Ile Phe Lys Thr Leu Gln Glu Glu Arg Arg Leu Arg Glu Ala
 565 570 575 580
 Ala Met Arg Ala Lys Val Glu Lys Thr Ala Leu Leu Lys Thr Glu Thr
 585 590 595
 55 Lys Glu Arg Thr Met Lys Ser Phe Leu Leu Ser Gln Lys His Val Val
 600 605 610

Tyr Thr Glu Pro Leu Asp Ile Gln Ala Gly Ser Ser Val Thr Val Tyr
 615 620 625
 5 Tyr Asn Pro Ala Asn Thr Val Leu Asn Gly Lys Pro Glu Ile Trp Phe
 630 635 640
 Arg Cys Ser Phe Asn Arg Trp Thr His Arg Leu Gly Pro Leu Pro Pro
 645 650 655 660
 10 Gln Lys Met Ser Pro Ala Glu Asn Gly Thr His Val Arg Ala Thr Val
 665 670 675
 Lys Val Pro Leu Asp Ala Tyr Met Met Asp Phe Val Phe Ser Glu Arg
 680 685 690
 15 Glu Asp Gly Gly Ile Phe Asp Asn Lys Ser Gly Met Asp Tyr His Ile
 695 700 705
 Pro Val Phe Gly Gly Val Ala Lys Glu Pro Pro Met His Ile Val His
 710 715 720
 20 Ile Ala Val Glu Met Ala Pro Ile Ala Lys Val Gly Gly Leu Gly Asp
 725 730 735 740
 25 Val Val Thr Ser Leu Ser Arg Ala Val Gln Asp Leu Asn His Asn Val
 745 750 755
 Asp Ile Ile Leu Pro Lys Tyr Asp Cys Leu Lys Met Asn Asn Val Lys
 760 765 770
 30 Asp Phe Arg Phe His Lys Asn Tyr Phe Trp Gly Gly Thr Glu Ile Lys
 775 780 785
 Val Trp Phe Gly Lys Val Glu Gly Leu Ser Val Tyr Phe Leu Glu Pro
 790 795 800
 35 Gln Asn Gly Leu Phe Ser Lys Gly Cys Val Tyr Gly Cys Ser Asn Asp
 805 810 815 820
 40 Gly Glu Arg Phe Gly Phe Phe Cys His Ala Ala Leu Glu Phe Leu Leu
 825 830 835
 Gln Gly Gly Phe Ser Pro Asp Ile Ile His Cys His Asp Trp Ser Ser
 840 845 850
 45 Ala Pro Val Ala Trp Leu Phe Lys Glu Gln Tyr Thr His Tyr Gly Leu
 855 860 865
 Ser Lys Ser Arg Ile Val Phe Thr Ile His Asn Leu Glu Phe Gly Ala
 870 875 880
 50 Asp Leu Ile Gly Arg Ala Met Thr Asn Ala Asp Lys Ala Thr Thr Val
 885 890 895 900
 55 Ser Pro Thr Tyr Ser Gln Glu Val Ser Gly Asn Pro Val Ile Ala Pro
 905 910 915

His Leu His Lys Phe His Gly Ile Val Asn Gly Ile Asp Pro Asp Ile
 920 925 930
 5 Trp Asp Pro Leu Asn Asp Lys Phe Ile Pro Ile Pro Tyr Thr Ser Glu
 935 940 945
 Asn Val Val Glu Gly Lys Thr Ala Ala Lys Glu Ala Leu Gln Arg Lys
 950 955 960
 10 Leu Gly Leu Lys Gln Ala Asp Leu Pro Leu Val Gly Ile Ile Thr Arg
 965 970 975 980
 15 Leu Thr His Gln Lys Gly Ile His Leu Ile Lys His Ala Ile Trp Arg
 985 990 995
 Thr Leu Glu Arg Asn Gly Gln Val Val Leu Leu Gly Ser Ala Pro Asp
 1000 1005 1010
 20 Pro Arg Val Gln Asn Asn Phe Val Asn Leu Ala Asn Gln Leu His Ser
 1015 1020 1025
 25 Lys Tyr Asn Asp Arg Ala Arg Leu Cys Leu Thr Tyr Asp Glu Pro Leu
 1030 1035 1040
 Ser His Leu Ile Tyr Ala Gly Ala Asp Phe Ile Leu Val Pro Ser Ile
 1045 1050 1055 1060
 30 Phe Glu Pro Cys Gly Leu Thr Gln Leu Thr Ala Met Arg Tyr Gly Ser
 1065 1070 1075
 Ile Pro Val Val Arg Lys Thr Gly Gly Leu Tyr Asp Thr Val Phe Asp
 1080 1085 1090
 35 Val Asp His Asp Lys Glu Arg Ala Gln Gln Cys Gly Leu Glu Pro Asn
 1095 1100 1105
 40 Gly Phe Ser Phe Asp Gly Ala Asp Ala Gly Gly Val Asp Tyr Ala Leu
 1110 1115 1120
 Asn Arg Ala Leu Ser Ala Trp Tyr Asp Gly Arg Asp Trp Phe Asn Ser
 1125 1130 1135 1140
 45 Leu Cys Lys Gln Val Met Glu Gln Asp Trp Ser Trp Asn Arg Pro Ala
 1145 1150 1155
 50 Leu Asp Tyr Leu Glu Leu Tyr His Ala Ala Arg Lys Leu Glu *
 1160 1165 1170

55 Claims

1. Altered starch extracted from a transformed potato plant or the progeny thereof, which starch, as extracted, has a viscosity onset temperature as determined by differential scanning calorimetry which is reduced by at least 5°C

compared to starch extracted from equivalent, non-transformed plants.

2. Altered starch according to claim 1, wherein the viscosity onset temperature is reduced by at least 7°C compared to starch extracted from equivalent, non-transformed plants.
- 5 3. Altered starch extracted from a transformed potato plant or the progeny thereof, which starch, as extracted, has a viscosity onset temperature as determined by differential scanning calorimetry, of less than 60°C.
- 10 4. Altered starch according to claim 3 which, as extracted, has a viscosity onset temperature of less than 55°C.
- 15 5. Altered starch according to any one of claims 1-4 which, as extracted, has a reduced endotherm peak temperature (as extracted) as determined by differential scanning calorimetry compared to starch extracted from equivalent, non-transformed plants.
- 20 6. Altered starch according to any one of claims 1-5 which, as extracted, has an endotherm peak temperature as determined by differential scanning calorimetry which is reduced by at least 5°C compared to starch extracted from equivalent, non-transformed plants.
- 25 7. Altered starch according to any one of claims 1-6 which, as extracted, has an endotherm peak temperature as determined by differential scanning calorimetry, of less than 59°C.
8. Altered starch according to any one of the preceding claims, having a substantially normal amylose content.
9. A polypeptide obtainable from a soluble extract of potato tubers and having starch synthase activity, in substantially isolated form.
- 30 10. A polypeptide according to claim 9, having an apparent molecular weight, as judged by SDS-PAGE, in the range of 100-140 kDa, or a functional equivalent thereof.
- 35 11. A polypeptide according to claim 9 or 10, having an apparent molecular weight, as judged by SDS-PAGE, of 140 kDa.
12. A polypeptide according to claim 9 or 10, having an apparent molecular weight, as judged by SDS-PAGE, of 120 kDa.
- 40 13. A polypeptide according to claim 9 or 10, having an apparent molecular weight, as judged by SDS-PAGE, of 110 kDa.
14. A polypeptide according to claim 9 or 10, obtainable from developing tubers of *S. tuberosum* cultivar Désirée, having an apparent molecular weight, as judged by SDS-PAGE, of 100 kDa.
- 45 15. A polypeptide according to any one of claims 9-14, comprising the amino acid sequence shown in Figure 6.
16. A nucleic acid sequence comprising at least 200bp and exhibiting at least 80% sequence identity with the corresponding region of the DNA sequence shown in Figure 6, operably linked in the sense or antisense orientation to a promoter operable in a plant.
17. A sequence according to claim 16 comprising at least 300-600bp.
- 50 18. A sequence according to claim 16 or 17, exhibiting at least 85% sequence identity with the corresponding region of the DNA sequence shown in Figure 6.
19. A sequence according to any one of claims 16, 17 or 18 exhibiting at least 90% sequence identity with the corresponding region of the DNA sequence shown in Figure 6.
- 55 20. A sequence according to any one of claims 16-19, comprising a 5' and/or a 3' untranslated region.
21. A sequence according to any one of claims 16-20, encoding at least a portion of a polypeptide in accordance with

any one of claim 15.

22. A sequence according to any one of claims 16-21, excluding sequences disclosed in WO 96/15248.

5 23. A nucleic acid construct comprising the nucleic acid sequence of any one of claims 16-22.

24. A host cell into which has been introduced a nucleic acid sequence according to any one of claims 16-22.

10 25. A host cell according to claim 24, wherein the nucleic acid sequence is introduced in a construct according to claim 23.

26. A host cell according to claim 24 or 25, wherein the introduced sequence is integrated into the host cell genome.

15 27. A plant host cell according to any one of claims 24, 25 or 26.

28. A plant or part thereof, into which has been introduced a nucleic acid sequence according to any one of claims 16-22, or the progeny of such a plant or part thereof.

20 29. A plant or part thereof according to claim 28, wherein the plant is selected from the group consisting of: potato, tomato, rice, wheat, pea cassava, sweet potato, barley, oat and maize.

30. A plant according to claim 28 or 29, comprising starch in accordance with any one of claims 1-8.

25 31. Starch extracted from a plant according to claim 28 or 29.

32. Starch according to claim 31, having altered properties, as extracted, relative to starch extracted from equivalent but untransformed plants.

30 33. Starch according to claim 31 or 32, and in accordance with any one of claims 1-8.

35 34. A method of producing altered starch from transformed potato plants or their progeny, the method comprising extracting starch from a potato plant, at least the tubers of which comprise a nucleic acid sequence in accordance with any one of claims 16-22, said sequence having been artificially introduced into the potato plant or a predecessor thereof.

40

45

50

55




Fig. 1

Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig 6

EcoRI NotI

GAATTCGGGGCCCGCATAGTGTGTGATGAAGGAGAGAGATATTTACATCGGATGCTTATTTGATTCGTGGTGAACAAAGTGTACAAAGAACATTCCTTTTCTTTTTC
 CTTAAGCGGGGGCTATACACACACTACTTCCTTCTCTATAAAGTGACCCACAAGATAACAGACACCACCTGCTTCAAAATGTTTCTTAAGGAAAAAAGAAAAAGG 120

NcoI

TTGGTTCTGTGTGGTCAAGCATGGATGTGCAATTCACATGCAATAGATCAATGAGTTCACAAAGTGTGCAATGCAATAACCCACCTCAAGAACAACCTATTCCTGGTGTGTCTC
 AACCAAGAACACACCCAGTGGTACCTACAAGTAAAGTGAAGTATGTAACCAACGTGTACAGAGGTACCTTATTTGGTGGAGTCTATTTGGATAAGAACCCAAAGAGAG 240

M D V F F P L H R D L S C T S V S N A I T - L K I K P L L S F V S

XmnI

TTATGGAAGCACAAAGTCTATCACTACAATCTCTTCATGGAGGAAGGATGGAATGGTACTGGGGTTCCTTTTCAATTTGTCCAAATTTCTCGGGAAGAACCGGAGAAAGTTTCAAC
 AGTACCTTTGGTGTGAGATAGTCACTGTAGAGGAAGTACCTCTCTTACCTTACCAATGACCCCAAGTAAAGGTAACACGCTTAAAGAGGCTTCTTCTGCTCTTTTCAAGTGTG 360

H S T T S L S V D S S S W R K D G M V L G V S F S I C A H F S G R R R R K V S T

ScaI

TCTAGGAGTCAAGGCTCTTCACTTAAGCGGTTTGTGCAAGGAAGCCCTCAGGGATGAGCAGCAAGAAAGGTTCAGAAAGCAATGGTGATAAAGAAAGTAAAGTACTTCAACATC
 AGGATCTTCAGTTCGAGAAGTGGATTCCTCAAAACAGGTTCTCTGGGAGTCTCTACTCTGGCTTCTTCAAGTCTTCTCTTACCACTATTTCTTTCATTTTCATGAAGTGTGAG 480

P R S D G S S P K G F V P R K P S G M S T O R K V O K S N G D K E S K S T S T S

TAAGAATCTCAAAATTCACACAGAGACGGTTGAAGCAAGAGTGAACATAGTGACGATGACACTAAAGGAGTGGTGAAGGACCAAGTTCTGGAGGATGAGGATGAATCAATCG
 ATTCTCTAGACTTAAAGGTTGGTCTTCTGCAACTTCTGTCACCTTATGATCACCTGTACTGTGATTTCTTCACTCTCTCTGTTTCAAGAGCTTCTTACTCTTACTTACTTACTT 600

K E S E I S N O K T V E A R V E T S O D D T K I V V R D H K F L E D E D E I N G

TTCTACTAAATCAATAGTATGTCTGCTGCTGATCTCAATTTGTGAAAGTGAAGAAAGTGGTGGTGTGATGACAAGGATGCTGTAAAGTAAACAAATCAAGAGATCGGAAGA
 AAGATGATTTAGTTATCATACAGTGGACAAAGACATAGTAGAGTAAACAACTTTCACCTCTTTTGACCACCCTACTGTTCTTACGACATTTCAATTTGTTAGTTCTCTAGCTTCT 720

S T K S I S H S P V R V S S D F V E S E E T G G D D K D A V K L N K S K R S E E

GAGTGGTTTTTATAATTGATTCTGTAAAGAGAACAAAGTGGATCTCAGGCGAAACTAAAGGAGTGAAGGAGGAGGATGCTGTGGGTACAAAGCTTTATGATATTTGAGGTGGA
 CTCACCAAAATTAAGTAAAGACATTTCTCTGTTTCACTTACAGTCCGCTTTGATTACGGTCACTGCTCTCTCTGCTACGACACCACTGTTTGAATACCTTAAGCTGCACT 840

S G F I I G S V I R E D S G S D G E T N A S S K G S H A V G T K L I E L C V D

TGTTGAGCCCAACAATTTGAAGAAATATCTTCAATGTGAATCAAAAGGACCTGTAGCAAGTAAAGTATTTCAAAATTAAGGCTAGTGTGCAACACACTGAAAGCAATGA
 ACAACTGGGTGTGTAACTTCTTTATTAAGACCTTACAACTTATGTTCTTGCACATGCTTATTGCAATACCTTAATGATTCGGATCACTACACCTTGTGTGACTTCTGTTACT 960

V E P C D L K E N N A G N V E Y K G P V A S K L L E I T K A S D V E H T E S N E

PstI

GATTGATGACTAGACACTAATAGTCTTTTAAATCAGATTTAATGAAGAGGATGAGCCATTAAGTTCACCAAGAGTGGAGACTGGAGATTTCTCTAAACTAAGATTGGAGATGGA
 CTAACTACTGAAATCTGTGATTATCAAAAGAAATTAAGTCTAAATTAATCTCTCTACCTGGTAACTGAGCTCTTGTCACTCTGACTCTAAGAGAGATTTGAATCTTAAGCTTCACT 1080

I D D L U T N S F F K S D L E E D E P L A A G T V E T G D S S L N L R L E N E

SnaBI

AGCAAACTACGTAGGCAAGCTATAGAAAGGCTTCCGAGGAAATTTATGCAAGGATCAGATTTATTTGTTTCCAGAGGTGTAAAGCTGATGAAGATGTCAGATATTTCTTAA
 TCGTTTGAATGATCTGTCGATATCTTTTGAAGGCTCTTTTAAATACGTTCTCTAGTCTAATAAAGCAAAAGGCTTCAAACTTTTGGATACCTTCAAGCTCTATAAAGAAAT 1200

A N L R R D A I E R L A E E N L L O G I R L F C F P E V V K P D E D V E I F L N

CAGAGGTCTTTTCACTTTGAAGAAAGTCTTATGTTTGTATTTGAGGAGCTTTTAAATGAGTGGGCTATAGGTCTTTTACACAGGCTAACGAGACTCATCTCAATGGAGATTGGTG
 GTCTCCAGAAAGGTGAAGCTTTTACTCAGACTACAGAACAAATACCTTCAAAATTAATTAAGGCTATCCAGAAATGATGTTCTGATTCAGCTCTGAGTACAGTTTACTTCAACAT 1320

R G L S T L K H E S D V L H D A F N E W R Y R S F T T R L T E H L N G D W H

GCTTTGCAAGATCCATGTTTCAAGGAAGCATACAGGCTGATTTGTGTTTTTAAATGGACAAGATGCTATGACAACAATGATGGAATGACTTCACTATACTGTGAAGGTTGGTA
 CAGAAGCTTCAAGTACAAGGTTCTTCTTCTATCTTCCACTTAAACACAAAAAATTAACCTGTTTCAAGATACTGTTGTTACTTACTTCACTTCACTTCACTTCACTTCACTTCACT 1440

S E K H V P K E A T R A D F V F F N G C D V Y D N N D G N D F S I T V K G G H

OFFICE OF THE DEPUTY ATTORNEY GENERAL

[illegible]

A A S E A O P A J A V E C A A K A K A V E R E L M / K A L Y R D I T A I F

~~GCGAATGAAATTAAATGGGAGGATAGGTCAGTTATACATAACAAAGGTCAGGTCCTCTCCCACTGCTAAGGACTGTGGATCGAGGGAGGATATAATATTTGGAGGATGCTTT
 TCTTCTGATTAATTAATACGGTCTCTCTGAGGATATGATATGCTTCTCAAGCTCAAGAGAGAGGGGTACGATTCTGAAACGCTATTTGCTCTCTATATTAATTAACCTCTTACGAAA~~

SSSEFKCEDK/APLRYNKS SGPLSHAKDL# : 10GVNN#KDC L

1920

5 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32

Sp1
GGATGGCATTCGTTATGATAAATCAGCCCAAGACTTCCATGCCATTGTCCCAACACCATCCGGAGGAATTATATTGGCTTGAGGAAGAACATCAGATCTTAAAGACACTCAGGA
CTACGCTAACCACTAATCTATTGTAAGTGGGGTTGTGAAGGTACGGTAACAGGGGTGGTGTAAAGGGCTCTTAATAACCCCAATCTCTCTGTAGCTAGAGAAATCTGTGAAGTCTT 2040

W A : A Y D N N H R O O F H A I V P N H I P E E L Y W V C E E H O I F K F L O O

GGAGAGAGGCGT*AGAGAGCGCGGT*CGCGCTAAGCTT*GAAAAACAGCGACTTCTGAAACGAAACGAAGGAAGCACTATGAAATCTATTTTCTGCTGAGAGCGATGAGTATA 2160
CGCTCTCGGAAATCTCTCGCGCGATAGCGGAGATCGCAACTTTTCTGCGTGAAGACCTTTGACCTTCTTCCCTCTTATATATGACAGATCTTCTCTGATCATAT

EXPLORE A MPAK: EKTALLK - ETKERTMK, SFL - SOKHV)

[illegible]

• E P L D I D A G S S S Y T V Y P N P A N T V L N G K P E • C R C S E N S U

CCGCGTGGGTCCATTGGCAGCTGCAAAAATGTCGCTGCTGAAAATGGCAGCTGCTGAGGCAAGCTGTGAAGTTCCTATGGCATGCAATATGATGGAATTTGTATTTGGCAGAGAGA 240
GGCGAGCGTACGGTAAAGGTGGAGCTCTTTACAGCGGAGCGCTTTTACGGCTGTATGATTTCTGTTGACACTTCGAAAGGTAACTCTGCTATATATGACGTAACATAAAACATAAAAGGCTGTGCT

P L C P L P O K H S P A E N G T H . R A T V K V P L C . M H C V F F D E P

ATATGGTGGGATTTTTCACATAAAGACGCGAAATGCACATATCCATACCTGTGTGTTGGAGGAGTCGTAAAGAACCTCCATGCAATCTGCAATATGCTGTATGCTGTGCGAATATGCAACCAATTCG 252
TCAACGACCTCAAAAAGTGTATTCTGGCTTACCTGTATGCTATATGGACAGCAACCTCTCTACCGCAATTCGTTGGAGGATACGCTAAAGATATTAACACACAGCTTACCTGTGCTAAAG

С. С. ГИФОНА С. И. ОУХИ. Р. В. С. ОУА КЕРПН. А. И. А. Е. Н. А. О.

[illegible]

K . S G L C D V S S R A F Q D H H N D I L P X O G L K P T

~~GTTGGGCTTCACAAAAAGTACTTTGGCGGTGGCACTGAATAAAGATATGGTTTGAAAGGTAAGAATGATGGC
ATTGTTTAGGCTTTTGTATTAATAGCAATCATTTATTTTATGCAAAATTTTCTAACCTCCAGAGAGCTAGATAAAGAGCTTGGGTAATAAAATCTTGGGCAC~~

Polymer Bulletin 1987, Vol. 10, No. 1, pp. 61-66
© VCH Verlagsgesellschaft mbH, D-6940 Weinheim, FRG

CGTCTATGCTTG74GCAACGA*GGTGAACGATTTGGTTCTCTATACGGGGGTTGGAGTTCTTCTGCAAGGTGGATTACTCCGA*ATCATTCATTGCCA*GATTGCTCTAGTGC
GCGATACCAACATCTGTGCTACCACTTGTAAACCAAGGAACAGACTGCCCGGAACCTCAAGAAGACGTTCCACCTAAATCAGGCTATAGTAAGTACGGTACTAACAGATCAGG 2880

V * S C S N D G E R F J F F C H A A L E F L L J G G F S P D I I H C W D W S S A

TCCTCTTGGTTGGCTCTTTAAGGAACAAATACGCACTATGGCTTAAGCAAACTCTGTATAGCTCTTCACGATACATAATCTTGAATTTGGGGCAGATCTCATTTGGGAGAGCAATGAC"AA
AGGACHAAGGACCGAGAAATTCCTTGTATATGTGTGATACCAATATGGCTTAGAGCATATAGGAAGTGGTATGTATTAGAAGCTTAAGAGCCCTCTAGAGTAACCCGTCTCTCTTACGAT" 3000

P V A W L F K E G I T H Y G L S K S R I V F T H N L E F G A D L I G R A M T N

CGCAGCAAGGTCACACAGCTTCCACCACTTACTCACAGAGGTTGTCTGGAACCTCTAATTCGCCCTCACCTTCACAGTTCACATGGTATAGTGAATGGGATGACCCAGATATTC
GGCTCTGTTTCGATGTCTGAAAGTGGTGAATGAGTGTCTCACAGACCTTGGGACATTAACGGGGAGTGGAGTGTTCACAGGTACCATATCACTTACCTTAACCTGGCTCTATAAAC

A C K A T T V S P T V S Q E V S G N P V I A P H L K F H S : V N G I D P J I Y

3240
G G A T C C T T T A A A C S A T A G T T C A T T C C G A T T C C G A C A T T C A G A A A A C G T G T T A G G C C A A A A C A G C A G C C A A G G A A G C T T T G C A G C G A A A C T T G G A C G A A A C A G G C T G A C T T C C
C C T A G G A A T T G C T A T C A A G T A A G C C T A A G C A T G T G A A G C T T T T G C A A C A C T C C C T T T T G T G T G G T C C T T T G C A A C G T G C G T T T G A A C C G A C T T G T C G A C T G S A A G G
O P L N D K K F I P I P Y T S E M V V F G K T A A K F A I D R Y I L L K O A D L R

TTTGTGAGGAATTATCACCCGCTTAACTCACCAGAAAGGAATCCACCTCAATAAACATGCTATTTCGCCACCTTGGAAACGGAACGACAGGTAGCTTGGCTGGTTCTGCTCTGTATCC 3360
AAACCACTCTTAATAGTGGCCGAATGAGTGGCTTCTCTTAGGTGGAGTAATTTGATGACATAAACCCCTGGAACTTGCCCTGCCGTGCATCAGAACGAACCAAGACGACGACTAGG

L V G I I T R L T H D K C I H L I K H A I W R T L E R N G G V V L L G S A P D P

T AGG GTAC AAA CAAT TT TGT TAAT TTGG CAA TCA ATG CACT CCA AAT ATA TAA TAC GCG CCA CAG CACT GT GTCT AAC ATAT GAC GAG CAC TT TCT ACC GTG ATAT GGT GGT GCG A
 A TCC ATG TTT GTT AAA CAAT TAA CCG TT AGT TAA CGT GAG GTT ATA TTA CTG CGC GGT GGT GAG ACAG ATT GT ATA CTG CT CGG TGA AGAG TGG ACT ATA TAC CAC CAG CACT 3680
 R Y V N N F V N L A V G L H S K Y N D R A R L C L T Y D E P L S H L I Y A G A D

1 T T A T T C T A G T T C C T T C A A T A T T T G A C C A T G T G G A C T A A C A C A C T T A C C G T A T A G A T A T G G T T C A A T C C A G T C G T C G C T A A A A C T G C A G C A C T T A T A G A T A C T A T T T C A T G T
 2 A A A A T A A G A T A A G G A A G T A T A A A C T C C G T A C A C C T G A T G C G T T G A A T C G G A T A C T C T A T A C C A A G T A A G G T C A G C A C C G A T T T T G A C C T C T G A A A T A C T A T S A C A T A A C T A C A
 3 F I L V P S I F E P C G L T Q L T A H R Y G S I P V V R K T S G L Y D T V F D V

TGACCATGACAAAGAGAGAGCACAACAGCTGTGGCTTGAACCAAAATGAAATACGCTTGTATGAGGAGATGCTGGCGGAGTGTATATGCTGTGAATAGAGCTCTCTGTGCTGGTACGA
 ACTGGTACTGTCTCTCTCGTGTGTGACACAGAACTTGGTTACCTAAGTGAAGAACTACCTGTCTACGACCGCTCAACTAATAACGAGACTATCTGACGAGAGACCAAGCAATGC
 D H G K E R A Q Q C G L E P N G F S F Q G A D A G C V D Y S I N R A I S A M Y D

GGTGGGATTGGTCAACTCTTTATGCAACGAGGTGATGCAACAAGATGGTCTTGGAAACGACCTGGCTTTGATTATTTGGAGCTTTACCATGCTGCTAGAAAGTAAAGTAGTAGT
ACCGGCTTAACCGAATGAGAAATACGTTGGTCCAGTAAGTGGTCTTAACGAGAACCTGGTGGGACGAGAACATAAAACCTGGAATGCTACGAGGATCTTCAATCTTATCATCA
G R D W F N S L C K D V M E Q D W S V N R P A L C Y L E L Y M A A R K L E

3960


ATAGCAGACAGCTCTCAAGTTCATTTTCATTTTTCGGGCAACATATGAAAGAGCTTAGCTCTTAAATAATCTCGCTATTTCATGATATTTCCTTTGGGAGAGAAATGAGAAATCAAG
TATCGTCTGTCAGGAATTCAGTAAAGTAAAGAAACCCCTTTTCATACTTTCGGAATCGGAGAAATATTACAGCCGGATAACTACTAATAACAAACCCCTTCTTACTCTTAGTTTC

GATGCAAAATAAAAAAAAAAAAAAAAAAAACTCGTGGCGAATTC 4127
CTACGTTTATTTTTTTTTTTTTTTTTTTTGGAGCACGGCTTAAG

A


pRAT4

Fig. 7



B

pPATRAT



2x 35S promoter

CaMV polyA

Polylinker


LB, RB Left and right border DNA

Patatin promoter

Nopaline synthase terminator

Neo Neomycin phosphotransferase

Pnos Nopaline synthase promoter



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) EP 0 779 363 A3

(12)

EUROPEAN PATENT APPLICATION

(88) Date of publication A3:
20.05.1998 Bulletin 1998/21

(51) Int Cl.⁶: C12N 15/82, C12N 15/54,
C12N 9/10, C08B 30/00,
C12N 5/10, A01H 5/00

(43) Date of publication A2:
18.06.1997 Bulletin 1997/25

(21) Application number: 96309004.8

(22) Date of filing: 11.12.1996

(84) Designated Contracting States:
AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE

- Marshall, Jacqueline
Warwick CV34 5SA (GB)
- Edwards, Elizabeth Ann
Norwich NR9 3DB (GB)
- Martin, Catherine Rosemary
Norwich NR15 1JW (GB)

(30) Priority: 12.12.1995 GB 9525353


(71) Applicant: National Starch and Chemical
Investment Holding Corporation
Wilmington, Delaware 19809 (US)

(74) Representative: Lipscombe, Martin John et al
Keith W Nash & Co,
Pearl Assurance House,
90-92 Regent Street
Cambridge CB2 1DP (GB)

(72) Inventors:
• Smith, Alison Mary
Norwich NR2 2BB (GB)

(54) Improvements in or relating to soluble starch synthase

(57) Disclosed is altered starch extracted from a transformed potato plant or the progeny thereof, which starch, as extracted, has a viscosity onset temperature as determined by differential scanning calorimetry which is reduced by at least 5°C compared to starch extracted from equivalent, non-transformed plants, together with a nucleic acid sequence comprising at least 200bp and exhibiting at least 80% sequence identity with the corresponding region of the DNA sequence shown in Figure 6, operably linked in the sense or anti-sense orientation to a promoter operable in a plant, and host cells and plants comprising the sequence.



EP 0 779 363 A3



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 96 30 9004

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.CI.6)
X, D	EDWARDS A ET AL.: "BIOCHEMICAL AND MOLECULAR CHARACTERIZATION OF A NOVEL STARCH SYNTHASE FROM POTATO TUBERS" PLANT JOURNAL, vol. 8, no. 2, August 1995, pages 283-294, XP002031375 * the whole document *	9	C12N15/82 C12N15/54 C12N9/10 C08B30/00 C12N5/10 A01H5/00
X	KOSSMANN J. ET AL.: "Transgenic plants as a tool to understand starch biosynthesis" PROGRESS IN BIOTECHNOLOGY, vol. 10, 23 - 26 April 1995, pages 271-278, XP002060102 * see p. 276/77 chapter 2.2.4. *	9-11	
A, D	WO 95 26407 A (NAT STARCH CHEM INVEST :COOKE DAVID (GB); GIDLEY MICHAEL JOHN (GB)) 5 October 1995 * the whole document *	1-34	
P, X, D	WO 96 15248 A (INST GENBIOLOGISCHE FORSCHUNG :KOSSMANN JENS (DE); SPRINGER FRANZI) 23 May 1996 * the whole document *	1-12, 15-21, 23-34	TECHNICAL FIELDS SEARCHED (Int.CI.6) C12N C08B A01H
P, X	MARSHALL J. ET AL.: "Identification of the major starch synthase in the soluble fraction of potato tubers" THE PLANT CELL, vol. 8, no. 7, July 1996, pages 1121-1135, XP002060103 * the whole document *	1-34	
The present search report has been drawn up for all claims			
Place of search		Date of completion of the search	Examiner
THE HAGUE		25 March 1998	Kania, T
CATEGORY OF CITED DOCUMENTS			
X particularly relevant if taken alone Y particularly relevant if combined with another document of the same category A technological background O non-written disclosure P intermediate document		T theory or principle underlying the invention E earlier patent document, but published on, or after, the filing date D document cited in the application L document cited for other reasons & member of the same patent family, corresponding document	

COU FORM 1503 03 02 (H04/201)